Plant Disease Reporter

Crops Research Division
Agricultural Research Service

UNITED STATES DEPARTMENT OF AGRICULTURE

VOLUME 51 NUMBER 13



1967 Index
(Issued December 1968)

The Plant Disease Reporter is published monthly as a service to American agriculture. The articles emphasize new developments in plant pathology, including disease incidence and severity, serious outbreaks and epidemics, investigative techniques, and control chemicals. The rapid issuance of such information permits plant pathologists to recommend the judicious use of pesticides or other appropriate control measures against impending disease epidemics. The Crops Research Division, in publishing this material, does not assume responsibility for the subject matter.

PLANT DISEASE REPORTER INDEX

Issued by

CROPS RESEARCH DIVISION

Plant Industry Station
Volume 51

Beltsville, Maryland
Number 13

AUTHOR INDEX

ACKERMAN, W. L., (715)
AFANASIEV, M. M., 269
ALEXANDER, L. J., (207)
ALLAN, R. E., 1041
ALLOW, JASSIM M., 809
AL-SOHAILY, I. A., (690)
ALTMAN, JACK, 86
ALVES, LEO M., 511
ARMSTRONG, G. M., 888
ARMSTRONG, JOANNE K., (888)
ASHOUR, W. E., (53)
ATKINS, I. M., (133), (536)
AUBÉ, CLAUDE, 573

BAGGA, H. S., 747, 855, 1055, 1058
BAGGETT, J. R., 661
BAKER, JAMES E., (1014)
BARKSDALE, T. H., 243
BARRY, J. ROBERT, 191
BASAK, WLADYSLAW, (3), 115, (117)
BEAN, GEORGE A., 839
BECKMAN, KENT M., (578)
BELL, A. A., 11
BELL, D. K., 986
BENIWAL, S. P. S., (861)
BERGESON, G. B., (78), (511)
BIRCHFIELD, W., 823
BLACK, L. L., 1038
BLODGETT, EARLE C., 302, (733), 893
BOCKHOLT, A. J., (777)
BOLLENBACHER, KATHARINA, 632
BOND, WILLIAM P., (108)
BOOTHROYD, C. W., (15)
BORECKI, Z., 3
BOYLE, ALICE M., (926)
BRADFUTE, O. E., (207)
BRAKKE, MYRON K., 1005
BRAMBL, ROBERT M., 381
BRENNAN, EILEEN, 850, 964
BRODIE, BILL B., 562
BROMFIELD, K. R., 248
BROOKS, T. L., 569
BROWN, G. ELDON, 95
BRUEHL, G. W., 810, 815
BUCKHOLTZ, W. F., (388)
BUGBEE, WILLIAM M., 760
BULLARD, R. G., (487)
BUMBIERIS, M., (473)
BURCHFIELD, E. G., (558)
BURTON, G. W., (562)
BUSHNELL, W. R., 447
BUTCHER, CLYDE L., 310
BUTTERFIELD, NORMAN W., (793)

CALVERT, OSCAR H., 377
CAMPBELL, FRANKLIN J., (793)
CAMPBELL, W. A., 929
CANTWELL, A. M., (493)
CARIS, MARLIN R., (476)
CARPENTER, J. B., 523, 525
CARROLL, VINCENT J., (345)
CATION, DONALD, 261
CAVENESS, FIELDS E., 33
CHAKRAVARTI, B. P., (764)
CHALKLEY, JAMES H., (235), (323), (419), (515)
CHAMBERS, ALBERT Y., (219), 771
CHANDLER, W. A., 257
CHAUBE, H. S., (1009), (1010)
CHAVEZ, H. B., 926
CHEE, K. H., 443
CHENEY, PHILIP W., (281), (509), (786)
CHENULU, V. V., (768)
CHIANG, M. S., (991)
CLARK, R. L., 1069
CLERK, G. C., 941
COBB, F. W., Jr., (856)
COLE, H., 40, 820
COLEMAN, O. H., (200)
CONES, WILLIAM L., 430
COOK, R. N., (839)
COPLIN, DAVID, (1066)
CORLEY, E. L., Jr., (482)
CORTÉS-MONLLOR, AMELIA, (739)
COVEY, RONALD P., Jr., 593
COYIER, DUANE L., 281, (509)
COYNE, D. P., 20, 25
CRANE, J. L., 112
CRÊTE, R., 991
CRITTENDEN, H. W., (112), (470)
CRONIN, EUNICE A., (766)
CROSSAN, D. F., (57), (158), (320), (495)
CULP, T. W., 856

DABBAH, ROGER, (296)
DAINES, ROBERT H., 349, (850)
DAKWA, J. T., (941)
DALE, J. L., 376, (381), (893)
DALELA, G. G., (800)
DANESH, DARIUSH, (595)
DAVIS, D. R., (72)

DAVIS, SPENCER H., Jr., (964)
DAVIS, TURNER S., 505
DEAN, C. E., 72
DEAN, L. L., (310)
DENNIS, GURDON K., (476)
DESCHÊNES, JACQUES, (573)
DEUTSCH, MIRIAM, 318
DIENER, URBAN L., 185
DIMOCK, A. W., 873
DINOOR, A., (120)
DOCAMPO, DELIA, (213), (396)
DOCHINGER, LEON S., (531), 900
DONALDSON, FRED S., Jr., 455
DONNELLY, E. D., (614)
DOWLER, WILLIAM M., 666
DUBIN, H. JESSE, 454
DuCHARME, E. P., 1031
DUICH, J., (40)
DUNLEAVY, J. M., 438, (872)
DURGAPAL, J. C., (928)

EASWARAN, K. S. S., 63
ECONOMIDES, C. V., (684)
EDMUNDS, J. E., 15
EL-HAIDERI, HAIDER, (371)
ENGLISH, HARLEY, 212, (213), (396)
EPPS, JAMES M., (771), 775
ESSER, R. P., 46
ETTINGER, G., (253)

FASSULIOTIS, GEORGE, 482, 557, 720
FAST, ROBERT, (20)
FATHI, SAAD M., (53)
FEDER, WILLIAM A., 793
FELDMAN, A. W., (675)
FELDMAN, J. M., (225)
FENWICK, HARRY S., (50), (591), 626
FIELDHOUSE, D. J., (320), (493), (497)
FINDLEY, W. R., (207)
FISHER, FRAN. E., 1070, 1071
FLEISHER, ZIPORA, 733
FLIEGEL, PETER, (155)
FLOWERS, R. A., 731, (866)
FOIL, R. RODNEY, 223
FORBES, IAN, (1001)
FORD, R. E., 388, 389
FRANK, Z. R., 414
FRAZIER, W. A., (661)
FREEMAN, K., (200)
FRENCH, D. W., (418)
FRENCH, WILLIAM J., 108
FRIDLUND, PAUL R., 42, (831), (1063)
FROSHEISER, F. I., 679
FULTON, N. D., (632)
FURR, J. R., (525)
FUTRELL, MAURICE C., 174

GALLETTA, G. J., (998)
GARNSEY, S. M., 410
GARREN, KENNETH H., 601
GARRETT, FRANK E., (185)
GATES, DONALD W., (385)
GHAFFAR, ABDUL, 743
GHOSH, A., (300), (1040)
GIBSON, KENNETH E., 976
GILL, D. L., 148, 971
GILLASPIE, A. G., Jr., 761
GILMER, R. M., 823
GILMOUR, J. W., 727
GOEL, L. B., 138
GOHEEN, A. C., (715)
GOLDEN, A. MORGAN, (164), (775), 974
GÖMEÇ, BELGIN, (893)
GOODING, G. V., Jr., 409
GORDON, D. T., (207)
GOSS, ROY L., (215)
GOULD, CHARLES J., 215
GREEN, RALPH J., Jr., 449
GREER, F. C., Jr., (868)
GRIFFIN, G. D., 651, 973
GROGAN, R. G., (841)
GROSS, HENRY L., 807, 941
GROVER, RAJENDRA K., 189
GUDAUSKAS, ROBERT T., 385
GUSTIN, R. D., (705)
GUTHRIE, JAMES W., 50, 591
GUTTER, Y., 58

HAGEDORN, D. J., 544
HANKS, R. W., 675
HANLIN, RICHARD T., 235, 323, 419, 515
HANSBROUGH, THOMAS, (223)
HANSEN, JOHN C., (298)
HANSEN, MELVIN C., 578
HARDING, PAUL R., Jr., 51, 781
HARJUNG, M. K., (629), (868)
HARLAN, D. P., 103
HARRISON, A. L., 441, 687
HARRISON, MONTY D., 9, (298)
HART, JOHN H., (230), 476
HART, W. H., (549), (794), 978
HARVEY, A. E., 1012
HAWKSWORTH, FRANK G., (791), 1049
HEALD, C. M., 581, (736), 905, 1035
HEHN, E. R., (702)
HELTON, A. W., 340, 655, (834)
HENDRIX, F. F., Jr., (929)
HENDRIX, J. WALTER, (911), (1074)
HEPTING, G. H., (4)
HERNANDEZ, TRAVIS P., (271)
HEWITT, W. B., (715)
HIGGINS, J. J., (164)
HOBBS, C. D., (777)
HOCKING, D., 276
HOLCOMB, G. E., 726
HOLLIS, JOHN P., 167
HOLTON, C. S., 844, 846
HRUSCHKA, HOWARD W., 1014
HURSH, JOHN S., (30)
HYRE, R. A., 253

ISELY, DUANE, (1017)
IVEY, HENRY, (148)

JACKSON, CURTIS R., 461
JENKINS, ANNA E., 785
JENKINS, L., (103)
JENSEN, HAROLD J., 98
JHOOTY, J. S., 1079
JINDAL, J. K., (928)
JOHNSON, LEANDER F., 219
JOHNSON, LOY C., (527)
JOLY, PATRICK, 296
JONES, JOHN PAUL, 645
JONES, J. W., (410)
JONES, L. G., (271)
JOSHI, L. M., (138), (646)
JOSHI, M. M., (1010)

KABLE, PHILLIP F., 155
KAHN, R. P., 715
KAISER, WALTER J., 595
KATCHO, ZUHAIR A., (809)
KATSANOS, R. A., 957
KAUFMAN, J., 696
KEIL, HARRY L, 345, 753
KENDRICK, E. L., (287)
KERR, ERIC D., 637
KHANNA, R. N., (1009), (1010)
KHARE, M. N., (353)
KHATRI, H. L., 768
KIECKHEFER, ROBERT W., 981
KILPATRICK, R. A., 133, 536, 539, (619), (826), 904
KIM, S. H., 497
KIMBLE, K. A., (841)
KIRKPATRICK, HUGH C., 786
KLUDY, DONALD H., (169)
KNUTSON, KENNETH W., (298)
KOCHAN, W. J., (340), (655)
KOIKE, H., 333, 491
KONSLER, T. R., (431)
KOTHARI, K. L., 686
KREBILL, R. G., 143
KREITLOW, K. W., 619
KRSTIC, MIHAILO M., 669
KRUGMAN, STANLEY L., (671)
KULIK, MARTIN M., (296)
KUNTZ, JAMES E., (160)
KYROU, N. C., 75, 936

LAFERRIERE, L., (310)
LAMBE, R. C., (388), (389), (438), 872
LAMOUREAUX, CLIFFORD H., (89)
LANDIS, W. RONALD, 230
LARSEN, M. J., (829)
LAUB, C. A., 659
LAUT, JOHN G., 899
LAWSON, R. H., 723, 952
LAZAR, A. C., 831, 1063
LEMBRIGHT, H. W., (331)
LENZ, J. V., (549), (978)
LEONE, IDA A., (850)
LEWIS, G. D., (709)
LIM, T. M., (443)
LINDNER, R. C., (786)
LITTRELL, R. H., 736
LIVINGSTON, CLARK H., (9), 298
LOEBENSTEIN, G., (318)
LORBEER, J. W., (696)
LORDELLO, LUIZ GONZAGA E., 791
LOWNSBERY, B. F., (841)
LUKE, H. H., (120), (915), (919), (923)
LUKE, W. J., (67)
LUKENS, R. J., 355, 752
LUTTRELL, E. S., (505)
LYDA, S. D., 331

MaCOMB, M. L., (705)
MacWITHEY, H. S., 83
MAGGENTI, A. R., 549, 794, (978)
MAI, W. F., (15), 398
MAIER, C. R., 487
MANNING, W. J., 158, (495)
MARTIN, NEIL E., 1074
MARTIN, W. J., (191), 271
MARTINEZ, A. L., 692
MATHENY, W. H., 169
MATHRE, D. E., 864
MATHUR, B. L., 629
MATHUR, R. L., (629), (800)
MATHUR, S. B., (138), 649
MAVRODINEAU, SIMONE, (1021)
McCAIN, ARTHUR H., 1051
McCARTNEY, WILLIAM O., 278
McCORNACK, A. A., (95)
McDANIEL, M. C., (71)
McDANIEL, M. E., (133)
McGLOHON, NORMAN E., (750)
McGREW, J. R., 288
McINTOSH, T. H., (926)
McINTYRE, GARY A., 608
McKEE, GUY W., (820)
McLEAN, D. M., 885
MERKLE, O. G., (536)
MERRIFIELD, R. G., (223)
METZGER, R. J., 287
MHAISKAR, V. G., 480
MICHEL, L. J., (130)
MIELKE, J. L., 306
MILHOLLAND, R. D., 998
MILLER, J. D., (120)
MILLER, P. M., 202, 609, 712
MILLER, V. L., (215)
MILLIKAN, D. F., (3), (115), (117)
MINTON, NORMAN A., 148, 614, 1001
MINZ, G., (373)
MISHRA, M. D., (300), (1040)
MOBLEY, CAROLE L., (539)
MOHAMED, HOSNI A., 53
MOLLER, W. J., (212), (213), (396)
MONROE, R. L., (715)
MOODY, A. R., 826

MOORE, LAURENCE D., 641
MOREY, D. D., (120)
MORTON, D. J., (158), 493, 495, (497)
MOTSINGER, RALPH E., (726), (823)
MUELLER, K. E., (595)
MUELLER, WALTER C., 1053
MUIR, JOHN A., 798
MULLIN, R. S., (1066)
MURPHY, H. C., 120, (391), (947)

NATOUR, R. M., 371
NAUER, E. M., (715)
NAULT, L. R., (207)
NEELY, DAN, 511
NEGULESCU, Fl., (358), (362)
NELSON, R. R., (110)
NEMEC, STANLEY, Jr., 283
NENE, Y. L., (822)
NOME, S. F., (212), (213), 396
NORTON, DON C., 1017
NOUR-ELDIN, F., (416)

O'BANNON, JOHN H., 995
ORELLANA, R. G., 182
ORR, CALVIN C., 29
OTTA, J. D., (864)
OU, S. H., (877)

PACUMBABA, R. P., 405
PALM, EINAR W., (377)
PAPASOLOMONTOS, A., 684
PAPAVIZAS, GEORGE C., 125
PAPPELIS, A. J., (957)
PAPPO, S., (733)
PARISH, C. L., (281), (509)
PARKER, K. G., (155), (398)
PARMETER, J. R., Jr., 856
PARRIS, G. K., 552, 805
PATHAK, V. N., 744
PATTERSON, SISTER MARY THOMASINE, 78
PÉREZ, J. ENRIQUE, 739
PERKINS, A. T., (40)
PERRY, V. G., (569), 1066
PETERS, DON, (527)
PETERSEN, D. H., (666), 750
PETERSON, GLENN W., 700
PETERSON, J. L., 436, 741
PETERSON, R. S., (306), 766
PHELPS, WILLIAM R., 160
PIFER, W. A., (715)
PINCKARD, J. A., 67
PINE, T. S., 64
PIRONE, THOMAS P., 108
PITRE, H. N., 1044
PONTIS, R. E., 225
POTTER, H. S., (622)
POUND, G. S., (136)
POWELL, DWIGHT, 351, 353, 605
POWERS, H. R., 4
PUNTER, DAVID, 357
PURCIFULL, D. E., 502
PURDY, LAURENCE H., 94, (1041)

QUALSET, C. O. 908
QUICK, CLARENCE R., 89

RAABE, ROBERT D., 390
RABBITT, A. E., (839)
RĂDULESCU, E., (358), (362)
RAHMAN, M. U., 758
RANGARAJAN, M., 764
RANNEY, C. D., 558
RAO, A. S., (1005)
RAO, Y. P., (928)
RATCLIFFE, T. J., (971)
RAU, G. J., (557)
RAYCHAUDHURI, S. P., 300, 1040
REED, HOWARD E., (219)
REEVES, E. L., (509)
REXRODE, C. O., 755
REYES, LUCAS, (120)
RHOADS, LARRY, (61)
RICH, AVERY E., 291, 293, (826)
RICH, SAUL, (712)
RIVERA, C. T., 877
ROBERTS, BRUCE R., 803
ROBISON, G. D., (331)
ROCHOW, W. F., 195
ROGERS, W. E., (491)
ROISTACHER, C. N., (715)
ROSS, ELDON W., 38
ROSS, MERRILL, (86)
ROTHMAN, PAUL G., 30, 354
ROWELL, J. B., 336, (447)

SAETTLER, ALFRED W., 622
SANKHLA, H. C., 800
SCHADE, JOHN E., (51)
SCHEFFER, R. P., 110
SCHIEBER, EUGENIO, 44, 265, 267
SCHNATHORST, W. C., 1066
SCHOENEWEISS, DONALD F., 933
SCHREIBER, LAWRENCE R., 531, (533)
SCHUSTER, M. L., (20), (25)
SEATON, G. A., (715)
SECHLER, DALE T., (120), 915, 919, 923
SEGALL, R. H., 151
SELF, R. L., (1035)
SHEPARD, J. F., (502)
SIGH, AMAR, (649)
SILVERBORG, S. B., (108), 829
SIMONS, M. D., 130, 947
SIMPSON, W. R., 540, 542, (626)
SINCLAIR, W. A., 1077
SINGH, B., (861)
SINGH, H. G., (800)
SINGH, R. S., 861, 1009, 1010
SINHA, S. K., 882
SIROHI, S. S., (189)
SIRRY, A. R., (53)
SLACK, D. A., (1038)
SLEETH, BAILEY, 138
SMART, GROVER C., Jr., (1066)
SMERLIS, E., 584, 678

SMILEY, J. H., (731), 866
SMITH, RICHARD S., Jr., 671
SMITH, WILSON L., Jr. (1014)
SMOOT, JOHN J., (95), 700
SOUTHARDS, CARROLL J., 455
SPAULDING, D. W., 225
SPECHT, C. H., (1021)
SRIVASTAVA, D. N., 928
SRIVASTAVA, H. S. LAL, (950)
STALEY, JOHN M., 791
STALL, R. E., (659)
STAVELY, J. R., 470
STEELE, A. E., 171
STEEPY, T. L., 709
STEGALL, W. A., Jr., (4)
STEWART, D. M., 358, 362
STEWART, VERN R., 702
STIPES, R. JAY, 533
STOKES, D. E., 508, 988
STOKES, G. W., (731)
STONER, WARREN N., 705, (981)
STRETCH, A. W., 401
STRICH-HARARI, DINA, 373
STRIDER, D. L., 431
SUBRAMANIAN, S., (758)
SUMNER, DONALD R., 457
SUTHERLAND, JACK R., 91, 367
SZTEJNBERG, A., 967

TAINTER, F. H., 418
TAMIMI, SADI A., 690
TANDON, I. N., 950
TAPPAN, W. B., (915)
TARJAN, A. C., 317, 1024
TAYLOR, A. L., (995)
TAYLOR, DONALD P., (974)
TAYLOR, G. S., (609)
TÉLIZ, DANIEL, 841
TODD, F. A., (409)
TOLER, ROBERT W., (108), 777
TOOLE, E. RICHARD, 500, 600
TROUTMAN, J. L., (856)
TRUJILLO, E. E., (1051)
TU, J. C., 682, 911
TUCKER, M. C., (71), 893

VAARTAJA, O., 473
VALENTA, V., 416
van der ZWET, T., (753)
VARNEY, E. H., (709)
VERMA, A. C., (686)
VERMA, S. C., (950)
VIDAVER, ANNE K., (396)

WADE, EARL K., (427)
WAHL, I., (120), (967)
WALKER, J. T., 1021
WALKEY, D. G. A., 883
WALLACE, J. M., (692), (715)
WALLIN, JACK R., 527
WALLNER, WILLIAM E., (476)
WALPER, SONIA, (454)
WALTER, RICHARD G., (9)
WASTIE, R. L., (443)
WATERWORTH, H. E., 164
WEATHERS, L. G., 629, 868
WEBSTER, ORRIN J., (174)
WEIDENSAUL, T. C., (807)
WEIHING, J. L., 396
WELLS, HOMER D., 468, (905), (1001)
WICKER, ED F., 311
WIEBEL, FREDERICK J., Jr., 57, 320
WILLIAMS, L. E., 207
WILLIAMS, PAUL H., 136, 427, 566
WILLIAMS, R. E., 834
WILLS, W. H., (641)
WILSON, CHARLES L., 71, 739
WINSTON, J. R., (700)
WINTERS, H. F., (715)
WOLTZ, S. S., (645)

YAMSONRAT, SUEBSAK, 960
YARWOOD, C. E., 178, 945
YASSIN, ABDEL MAGEED, 30, 1028
YOUNG, P. A., 314, 464
YOUNT, WILLIAM L., 61

ZAK, B., 264
ZAWADZKA, BARBARA, (3), (115), 117
ZENTMYER, GEORGE A., (267)
ZILLINSKY, F. J., 391
ZIMMER, D. E., 586, 589
ZUMMO, N., 200

SUBJECT INDEX

Abaca: propagation of virus-free suckers of mosaic-infected abaca corms, 405
Abies concolor: fall soil fumigation effective in control of charcoal root disease in white fir and ponderosa pine nursery seedlings, 671
Abrasives: use of Celite as abrasive, increased virus transmission in mechanical inoculations, 945
Aceria tulipae: 207
Aerobacter cloaceae: 151
Africa: 416
Agar (see Cultures)
Agropyron smithii: non-rosaceous host of Phragmidium (Uredinales), 766
Air pollution: characterization of the plant damage in New Jersey, 850; damage to Austrian pine in New Jersey following period of high air pollution, 964

Alabama: 148; 385
Alfalfa: first survey in Quebec of fungi associated with crown and root rots of alfalfa and red clover, 573; inoculation of seed, seedlings or mature plants with suspension of nematodes as techniques for screening for resistance to stem nematodes, 651; physiogenic leaf and stem spot, 619; Phytophthora root rot, caused stand depletion in Minnesota, 679; stem nematode, liquid formulations of cynem and dimethoate gave good control, 973
Almond (see Prunus)
Alternaria spp.: key to the most common species of the genus Alternaria, 296
--- solani: 9
--- tenuis: 151
--- triticina: 950
Amendments: alfalfa hay, oat straw, lespedeza hay and flax hay reduced root-knot infection of tomatoes in field experiments in Tennessee, 219; cruciferous leaves and stems suppressed root rot in greenhouse treatments for control of Aphanomyces root rot of peas, 125; sawdust + balanced NPK fertilizers applied to soil suppressed root knot of okra and tomato and increased yield in India, 861; soybean meal and certain plant-derived oil-amended soils showed reduction in number of lesion nematodes, 1021
Announcements: 319; 322
Anthracnose: of poplar, first North American record of pathogen in Oklahoma, 785
---: of rye, major disease limiting seed production in Florida, 923
---: seedling: of cotton, susceptibility of 24 varieties of Gossypium hirsutum compared to Nanking variety of G. arboreum, 632
Antibiotics: cycloheximide derivatives on loblolly pines infected with fusiform rust, 223; sprays fail to affect fusiform rust galls on loblolly pine, 505; cycloheximide thiosemicarbazone on Cytospora canker disease of 'Italian' prune trees, evaluated for first and second year effects, 655
---: Oligomycin treatments of oak wilt in northern pin oaks, 160
---: Penicillium rubrum strain possesses fungistatic and fungicidal properties toward several plant pathogens and wood colonizing fungi, 669
---: phytoactin applied to western white pine seedlings did not affect normal mycorrhizae-root consociations, 1012
---: streptomycin used with regular bean seed treatments indicated no significant control of bean halo blight, 544; night applications of streptomycin more effective than daytime applications in preventing fire blight disease, 605
---: Terramycin: for control of peach bacterial spot, 345; looks promising as control for peach bacterial spot infection, on fruit and foliage alike, in Illinois, 351; Terramycin, 132 ppm and streptomycin, 150 ppm gave superior control of bacterial spot on peach fruit and foliage alike, 349
Aphelenchoides fragariae: 988
Aphelenchus avenae: 367
Apple: canker: Jonathan apple new host, 593; first report of fruit rot phase of European canker disease of apple in European canker orchards in California, 278
--- crab: trunk canker of ornamental crabapples, isolation of Physalospora obtusa and three other microorganisms from cankered trees, 230
---: fire blight: first authentic report of disease on continental Europe, 3; night applications of streptomycin more effective than daytime applications for preventing fire blight disease, 605
---: flat apple disease found to be graft transmissible on Red Delicious apple trees, 509
---: transmissible "red ring" disease described on Red Delicious apples, 281
---: sodium hypochlorite as disinfectant of pruning tools for fire blight control on Jonathan apple twigs, 753
---: stem cavity russet, occurrence and amount on Golden Delicious apples, 302
---: viruses: influence of dapple apple and stem pitting viruses on tree growth and fruit yields and storage life, 293; stem pitting virus, most natural transmission could be by root grafting as suggested by pattern of spread, 291
Apricot (see also Prunus): rust, caused by Tranzschelia discolor, in Chile for first time, 1966, 212; silverleaf, caused by Stereum purpureum, previously unreported in Chile, 212
Arceuthobium campylopodum: 856
--- --- f. campylopodum: 311
--- pusillum: eastern dwarf mistletoe, on jack pine, 899
--- vaginatum cryptopodum: 1049

Argentina: 225
Arizona: 1049
Arkansas: 71; 376; 893; 1038
Armillaria mellea: 390
Ash: green, root rot caused by Corticium galactinum developed in three of 10 seedlings when pathogen was placed in contact with tap roots, 500
Asparagus: ditch bank weed host for asparagus rust in south Idaho, 542
Aspergillus spp.: 461; 523
Australia: 473
Azalea: 10-year summary of plant-parasitic nematodes associated with azalea in Florida, 14 genera and 20 species reported for first time, 46

Bacteria: carbohydrate utilization by Xanthomonas vesicatoria, influence of length of time in culture upon, 57; viability of microorganisms prolonged by drying cultures to desiccation level and storing in sealed jars at 10-13°C temperature, 747
---: plant-pathogenic, use of sheep to produce large quantities of high titer antiserum against, 50
Bacterial blight (see Blight)
--- canker: in peach trees, transmission by bud propagation, 666; of tomato, found for first time in Chile, 396
--- leaf blotch: of sorghum, seed transmission, in south India, 63
--- --- streak: of rice, in India, 928
--- necrosis: of tomato, increase in incidence caused by holding mature-green tomatoes at chilling temperatures after harvest, 151
--- rot (see Rot)
--- spot: leaf spot and stem rot of geraniums in Pennsylvania, 61
--- ---: of peach, antibiotics Terramycin and streptomycin gave superior control of bacterial spot on fruit and foliage alike in New Jersey, 349; Terramycin gave better bacterial spot control in 2 years of field experiments than any other material in Maryland, 345; Terramycin may be promising as control for peach bacterial spot infection in Illinois, 351
--- ---: of pepper, effect of growth-regulating and other compounds on, 320
--- ---: of tomato, an evaluation of nightshade and ground cherry as suspects of Xanthomonas vesicatoria, 659; several species, varieties and plant introductions tested for reaction of foliage to Florida and Nebraska isolates of Xanthomonas vesicatoria, 25
--- stalk rot: of corn, in India, 764

Barley: effects of temperature, seed treatment, and soil fumigation on stand reduction of seedlings, 919
---: root-knot nematode, first report of occurrence in Illinois, 974
---: yellow dwarf virus: aluminum foil not effective in protecting winter oats from aphid vecotrs, 354; report of predominating strains of BYDV in New York showing changes during 10-year period, 195
Bean: circular leaf spot of french beans caused by new virulent form of Phaeoisariopsis griseola, 276
---: downy mildew, effect of temperature, time, and light on the quantitative production of sporangia, 253
---: halo blight: incidence in Idaho seed fields, 310; in pathogenicity tests all 17 Idaho isolates yielded a race 2 reaction on six bean varieties, 591; of snap bean, field and greenhouse tests of commercial varieties and breeding lines for sources of additional resistance to halo blight, 661; several bean varieties tested in early and late summer field plantings for reaction to races and strains of bacteria, 20; streptomycin used with regular bean seed treatments indicated no significant control of bean halo blight in Wisconsin, 544
---: root and hypocotyl rot of snapbean, depth of planting appears important factor in incidence of rot, 158
Bifusella crepidiformis: 791
Blackleg: of cabbage, occurrence of Phoma lingam on cabbage seed from Australia after treatment with hot water, 566
Black shank: of tobacco, calcium nutrition in relation to susceptibility of tobacco seedlings, 641; distribution of race 1 of Phytophthora parasitica in Kentucky and Tennessee, 731
Blackstem canker: on cottonwood, results indicate disease may be a symptomatic response of weakened host tissue to invasion by fungi, 933
Blast disease: of millet, first report of Piricularia setariae as a seedborne pathogen in Setaria italica, 138
--- ---: of rice, spread in small fields, 243
Blight: bacterial: fire: first authentic report of the disease on continental Europe, found on pear and apple trees, 3; sodium hypochlorite as disinfectant of pruning tools for control of, 753
---: ---: halo: and common blight of

bean, report on current status of chemical control of bacterial blights of dry field beans in Michigan, 622; bean varieties tested for sources of tolerance and reaction to races and strains of Pseudomonas phaseolicola, 20; of bean seed, incidence in Idaho, 310; on snap bean, sources of resistance in commercial varieties and breeding lines reviewed, 661; pathogenicity of Idaho isolates tested on six bean varieties, 591; streptomycin seed treatment for control of bean halo blight indicated no significant control in Wisconsin, 544
---: Colletotrichum, of dwarf mistletoe on Pinus ponderosa, effects a degree of biological control, 311
---: early, of potato, tetrachloroisophthalonitrile evaluated and compared with maneb for control under field conditions, 9
---: Fusarium: of turfgrass, effectiveness of chemical control of, 839; of zoysia in Arkansas, developed under abnormal environmental conditions, new disease caused by Fusarium spp., 376
---: late: of potato, relationship of rainfall and temperature to occurrence of disease in Greece, 11 years of data analyzed, 75; three methods of analyzing weather data in relation to occurrence of late blight of potato in Thessaloniki, Greece, 936
---: leaf, of wheat, effectively controlled in India by fungicidal sprays, 950
---: needle: of Douglas-fir seedlings in Chile, 454; of pine, distribution and significance in New Zealand, 727
---: sclerotial, of sweetpotato, controlled by 2,6-dichloro-4-nitroaniline, 191
---: sheath, of rice in Taiwan, 20 strains of Pellicularia sasakii isolated, 682
---: southern, its control on tomato and peanut, effective control on peanut production with 3-inch high beds, deep plowing, and PCNB treatment, 314
---: snow, of fir, pathogenicity of Phacidium abietis on several conifer species, 678
---: twig, new disease of Vinca rosea in India, 480
Blueberry: occurrence and control of Glomerella cingulata on highbush blueberry, 401; Phytophthora root rot, relative susceptibility of blueberry cultivars to Phytophthora cinnamomi, 998
Botrysphaeria ribis: 593
Botrytis allii: control of Botrytis neck rot in late harvested onions in New York by fungicidal dusts, 696
--- cinerea: 212
--- convoluta: 83
--- fabae: 595
Boxwood: Japanese, excellent control of root-knot nematodes with phosphorate nematocides, 1035
Brazil: 791
Brown rot: blossom blight of almonds, excellent control from DCNA, 578; of peach leaf blades, evidence of direct penetration of germinating conidia, 750
--- spot: of loblolly pine, field infection of pine seedlings with naturally produced inoculum, 552
Buffaloberry: leaf spot of seedlings controlled with captan or maneb, 700
Bunt: of wheat, new race of Tilletia caries (designated X-1) identified that attacks PI 192383, PI 116301, and M63-1223, 287

Cabbage (see Crucifers)
California: 278; 390; 671
Canada: 798
Canker: on Jonathan apple trees, first report of Botryosphaeria canker on apples from Washington, 593
---: on ornamental crabapple, an attempt to determine cause and extent of disease on Michigan State University campus, 230
---: bleeding, Phacidiopycnis pseudotsugae associated with, on hemlock, first report of fungus affecting hemlock naturally, 807
---: collar, of coffee trees in two new locations in Guatemala, 267
---: of cottonwood and hybrid poplar trees, first report of occurrence in Iowa, 900
---: Cytospora: of black cherry, excessive gum production in relation to cambium miner insect, 941; on sweet cherry, associated with winter injury, 155; on 'Italian' prune trees, evaluation of first and second year effects from treatment with cycloheximide thiosemicarbazone on old and young trees, 655
---: European: eye rot of apple, caused by Nectria galligena, first report of fruit rot phase in California, 278
---: Fusarium canker on paper mulberry, apparently a weak parasite attacking dormant trees or trees under environmental stress, 531
---: Scleroderris: of red pine in New York plantations, 108; on red pine, first confirmed record on seedling

stock in forest nurseries in Ontario, 357
Carbohydrate: utilization by Xanthomonas vesicatoria, influence of length of time in culture upon, 57
Carnation: tip burn on, 793
Celeriac (root celery): scab causes losses as high as 50% in Wisconsin, first report of scab on celeriac in U.S., 427
Celery: Fusarium wilt, caused wilt of celery and garden pea, 888
---: western celery mosaic virus identified in celery from commercial plantings in Florida, 502
Centrospora acerina: 971
Cephalosporium gregatum: 438
Ceratocystis fagacearum: 430; 755
--- minor: 669
--- (Ophiostoma) roboris: 739
--- ulmi: 71; 476; 511; 533; 803; 1077
Cercospora kikuchii: growth on various media, 112
--- personata: 687
Chemicals: several field tested for control of bacterial spot of pepper, 497; effect of SMDC on germination, plant growth and Fusarium wilt incidence of cumin in India, 629; excess of calcium carbonate in soil made iron mostly unavailable to peanut roots causing chlorosis, 464; organic spray materials reduced incidence of stylar-end rot of limes, chemicals may have served as source of nutrition, 1071; fluorochemical liquid FC-43 effective carrier for conidia of Erysiphe graminis or uredospores of Puccinia graminis to plant surfaces, 447
---: growth regulators, effect on bacterial spot of pepper, 320
---: hydrogen peroxide, effect of temperature and hydrogen peroxide on extraction of burrowing nematodes from citrus roots, 1024
---: sodium hypochlorite, as disinfectant of pruning tools for fire blight control, 753
Cherry (see also Prunus): preplant fumigation of planting sites by nematocides resulted in greater tree growth and decrease in nematode population, 398
---: canker: Cytospora canker of black cherry, and relation of cambium miner insect to excessive gum production, 941; associated with winter injury, 155
---: viruses: indexing of 47 varieties of cherry trees in New York resulted in isolation of apple chlorotic leaf spot virus (CLSV) and tobacco mosaic virus (TMV), 823; of sweet cherry, effect of temperature on symptom development of Lambert mottle of, 269
Chicory: Rhizopus rot of, new disease in India, 189
Chile: 212; 213; 396
Chocolate spot: of broadbean, with broadbean rust and bean yellow mosaic virus reached epidemic proportions in Iran 1966, 595
Chrysanthemum: Fusarium wilt of, effect of Meloidogyne hapla and Fusarium oxysporum on severity of, 736; three aspermy related viruses from chrysanthemum not identical to tomato aspermy virus, 723
Citrus spp.: Cladosporium leaf spot, not previously described on citrus in Florida, 1070
---: dieback, Aspergillus foetidus and Aspergillus sp. isolated from lesions on young nursery stock and top-worked citrus trees, 523
---: nematodes: annual population periodicity of Radopholus similis in Florida citrus groves, 1031; distribution of citrus nematode in Florida, 317; effect of temperature and hydrogen peroxide on extraction of burrowing nematodes, 1024; effective control on citrus seedlings using nematocidal bare-root dips, 995; pathogenicity of Pratylenchus brachyurus to citrus established by inoculating sour orange seedlings growing in sterile soil, 569
---: postharvest decay: biphenyl-resistant citrus green mold reported in Florida, 700; fruit rots, new method of applying biphenyl for control, 58; control of Penicillium mold on lemons, 781; Penicillium decay, of navel oranges, testing thiabendazole in resin-solvent and wax-emulsion coatings for control, 51; thiabendazole as a postharvest fungicide, 95
---: stylar-end rot of limes reduced by use of organic spray materials, 1071
---: viruses: 'Alemow' Citrus macrophylla, used as rootstock for lemons in California susceptible to cachexia virus, 525; cross-transmission studies with Citrus and Prunus viruses, 64; effect of rootstock on incidence of impietratura-diseased grapefruit fruits in Cyprus, 684
---: exocortis virus: of citron plants apparently spread by use of contaminated budding knife, 410; successful transmission of exocortis virus by dodder, grafting and mechanical methods, 868

---: leaf-mottle-yellows disease, transmitted by psyllid Diaphorina citri, may be virus responsible for severe tree losses in last 10 years in Philippines, 692
Cladosporium cucumerinum: 495
--- oxysporum: 1070
Clover: a physiogenic leaf and stem spot, 619; root rot, cutting height affected incidence of in greenhouse experiments, of minor importance in field trial, 826; survey of fungi associated with crown and root rots of alfalfa and red clover, first work of this kind in Quebec, 573
Clubroot: of crucifers, screening tests of crucifers for resistance to race 6A in organic soils of Quebec, 991
Coffee: collar canker, first report from two locations in Guatemala, 267; new Fusarium wilt of, in India, 758
Colletotrichum sp.: 480
--- gloeosporioides: 311; first report on dwarf mistletoe in Canada, 798
--- gossypii: 632
--- graminicola: 923; 957
--- orbiculare: 885
Comandra umbellata: alternate host of comandra rust of hard pines in the Cumberland Plateau in eastern Tennessee, 4
Conifers: organisms associated with root rots of, in South Australian nurseries, 473
Connecticut: 752
Corn: fungi carried by seed and their importance in causing diseases in the U.A.R., 7 genera isolated from 42 samples, 53; used as rotation crop with cucurbits for reducing cucurbit scab, 495
--- : bacterial stalk rot on, caused by Pseudomonas lapsa, in India, 764
---: downy mildew, two downy mildews of corn found to be internally seedborne in India, 1010
---: holcus leaf spot, epidemic in east-central Nebraska in 1966, 396
--- rot: influence of cultural practices on the incidence of stalk rot, 540; root and stalk rot, D-D not recommended for control, 15
--- smut: in greenhouse trials in-furrow treatment with PCNB resulted in complete control of head smut of corn, 626; PCNB applied as in-furrow treatment for control of head smut of corn, 626
---: viruses: occurrence of mechanically transmissible virus of corn and small grains in Ohio, 207; nonpersistent virus found infecting corn in South Dakota, 705; positive relationship between virus ratings and survival of plants with ears, regardless of specific virus or viruses involved, 377
---: corn stunt virus, and mechanically transmissible mosaic virus isolate found on Johnson grass, effect on dent corn hybrid in Mississippi, 1044
---: maize dwarf mosaic: additional data on distribution and properties in Alabama, 385; identification, distribution, and transmission of, 777
---: wheat streak mosaic, incidence in Iowa in 1966, 389
Corrections: 234; 896
Corticium fuciforme: 215
--- galactinum: 500
Corynebacterium michiganense: 396
Cotton: reduction in 1966 yield caused by diseases, 370; sources of resistance in the genus Gossypium to several soilborne pathogens, 864
---: bacterial blight, absence of contamination with Xanthomonas malvacearum in cotton gins 5 years after field eradication in California, 1066
---: boll rot: Pellicularia filamentosa found to be a primary cause in Louisiana, 67; storage of boll rot organisms on infected cotton bolls for prolonged periods, 855
---: Microxyphiella sp., apparently first report on, 743
---: nematodes, root-knot nematode damage observed in cotton after several years of continuous sorghum, 29
---: root rot: fumigants for control of, 331; greenhouse infection of cotton by Phymatotrichum omnivorum affected by temperature, 926
---: seedling disease: evaluation of granular formulations of soil fungicides for seedling disease control, 487; seed treatment, chloroneb used as overtreatment of mercury-treated cottonseed not so effective as captan-PCNB in Mississippi tests for seedling disease control, 558; seedling anthracnose, susceptibility of 24 varieties of Gossypium hirsutum compared to Nanking variety of G. arboreum, 632
---: Verticillium wilt, first report of isolation of Verticillium nigrescens from cotton, in Missouri, 760
Cottonwood: blackstem canker: susceptibility of weakened cottonwood stems to fungi associated with blackstem, 933; canker disease, first report of occurrence in Iowa, 900
Crabapple (see Apple)
Criconemoides sp.: 905

Cronartium comandrae: 4
--- conigenum: 44
--- fusiforme: 223
--- ribicola: 89
Crop rotation: for control of cucurbit scab, 1-year rotation with corn reduced scab incidence on cantaloup and cucumber fruit in Delaware, 495
Crown rust: of oats, determination of relative tolerance to Puccinia coronata avenae of experimental lines of oats, 947
Crucifers: blackleg, occurrence of Phoma lingam on cabbage seed from Australia after treatment with hot water, 566
---: clubroot, five cabbage selections and one variety and two kale varieties show some resistance to race 6A of Plasmodiophora brassicae in Quebec, 991
---: powdery mildew, first record of perithecial stage of the fungus found on Brassica campestris var. sarson and Brassica juncea in India, 800
---: White Spike radish, development of a Fusarium-resistant intermediate length type, 136
Cryptosporella viticola: 1038
Cucumber (see Cucurbits)
Cucurbits: five of 14 species of Cucumis resistant to root-knot nematode in greenhouse screening, 720; identity of powdery mildews of cucurbits in India, 1079
---: cantaloup, cantaloup and cucumber plantings showed marked reduction of cucurbit scab by a 1-year rotation with corn, 495
---: cucumber, trellised, varietal performance during serious outbreak of cucumber mosaic virus in North Carolina, 431
---: pumpkin, curly top virus found in Iowa in 1967, 1069
---: watermelon, interaction of race I and race II of Colletotrichum orbiculare on watermelon, 885
Cultures: growth of Cercospora kikuchii on various media, 112; influence of length of time in culture upon carbohydrate utilization by Xanthomonas vesicatoria, 57
Cumin: effect of SMDC on germination, plant growth, and Fusarium wilt incidence of, 629
Curvularia spp.: 11; 376
Cylindrocladium crotalariae: 986
--- scoparium: associated with extensive root damage and mortality in a 27-year-old yellow-poplar plantation, 38
Cylindrosporium spp.: 700
Cyprus: 684
Cytospora sp.: 155; 212; 340
--- chrysosperma: 933
--- cincta: 655; 834
--- leucostoma: 941

Dead arm disease: of grape, cane and leafspot phase of found in Arkansas and West Virginia, 1038
Decay: Penicillium, of navel oranges, testing thiabendazole in resin-solvent and wax-emulsion coatings for control, 51
Delmarva Peninsula: 493
Dieback: of citrus, association of Aspergillus spp. with, 523
Diplodia natalensis: 95; 744
Diseases: Index to genera of pathogens listed in the "Index of Plant Diseases in the United States"--4 parts, 235, 323, 419, 515; reduction in 1966 yield of cotton caused by, 370; of tobacco, in Kentucky in 1966, 866
Distribution: additional data on the distribution and properties of maize dwarf mosaic virus in Alabama, 385; distribution and prevalence of virulence for Victoria oats in Ustilago avenae in the United States, 846; of citrus nematode in Florida, 317; of pome fruit viruses in Poland, 117; of Helminthosporium victoriae, 110; of race 1 of Phytophthora parasitica nicotianae in Kentucky and Tennessee, 731; of stone fruit viruses in Poland, 115; of stunt disease of peanut in Virginia in 1966, 169; of Verticillium albo-atrum and V. dahliae in commercial potato-growing regions of Maine, 608
Ditylenchus dipsaci: 651; 973
Dodder: transmission of citrange stunt virus of citrus by dodder, 629; transmission of exocortis virus of citrus by, 868
Dollar spot: on bentgrass, varietal susceptibility to infection and influence of variety on fungicide effectiveness, 40
Dothistroma pini: 454; 727
Downy mildew: of corn, internally seedborne nature of two downy mildews on corn, 1010; of lima bean, effect of temperature, time, and light on the quantitative production of sporangia, 253; method for obtaining quick infection with Sclerospora species; 1009; of opium poppy, root applications of certain fungicides show partial control in India, 686
Dutch elm disease: control studies with captan and other fungicidal agents,

533; distribution in Arkansas in 1966, 71; in Illinois cities, 511; radiocarbon used for labeling conidia of Ceratocystis ulmi, 803; summer trimming of healthy trees attracted Scolytus multistriatus, first report of S. multistriatus attempting to form brood galleries in recently trimmed, healthy elm trees, 476; three carboxylic acids show no promise for prevention or therapy of disease, 1077
Dwarf mistletoe: bishop pine mistletoe seeds found to infect two Monterey pine seedlings in experimental testing, 856; Colletotrichum blight on, evaluation of biological control of dwarf mistletoe, 311; Colletotrichum gloeosporioides, first report in Canada, 798; first authentic record of Arceuthobium pusillum infection on jack pine in southeastern Manitoba, 899; on eastern larch in Minnesota, 418; on ponderosa pine, distribution on South Rim of Grand Canyon, Arizona, 1049

Elder: tobacco mosaic virus, first known report of natural infection of, 1053
Elm: Dutch: conidia of Ceratocystis ulmi successfully labeled with radiocarbon, 803; control studies with captan and other fungicidal agents showed no significant control, 533; distribution in Arkansas in 1966, 71; first report of S. multistriatus attempting to form brood galleries in recently trimmed, healthy trees in Michigan, 476; in Illinois cities, 511; three carboxylic acids on young elm trees show no promise for prevention or therapy of disease, 1077
Endothia parasitica: 669
England: 883
Epidemics: of broadbean crop in Iran in 1966, 595; holcus leaf spot on corn in east-central Nebraska, 396; of powdery mildew in winter oats in Stoneville, Mississippi, 30
Epidemiology: experimental initiation, with Puccinia graminis tritici, the minimum data that seem necessary for adequate prediction of field infection, 248; possibility that saprozoic nematodes could have a significant role in epidemiology of plant diseases, 98
Erwinia amylovora: 3; 605; 753
--- carotovora: 151
Erysiphe graminis: 447
--- --- avenae: 30; 391
--- --- tritici: 94
--- polygoni: 800
Eucalyptus: Pratylenchus brachyurus, first report on, 791
Euphorbia lagascae: Meloidogyne hapla and Sclerotium rolfsii of, 164
Europe: 3

Fabraea maculata: 257
Fern: newly reported hosts of Aphelenchoides fragariae in Florida, 508
Fig: mosaic virus, no nursery stock free of symptoms could be found in Turkey, 893
Fir (see Abies and Pseudotsuga)
Florida: 46; 72; 317; 502; 508; 700; 919; 923; 1031
Fomes pini: 829
Fungi: associated with Zoysia japonica in the Washington, D.C. area, 11; drying cultures to desiccation level and storing in sealed jars at 10-13°C temperature prolonged viability of microorganisms, 747; organisms isolated from stalk rot complex in Idaho, 540; seven genera isolated from 42 corn samples representing the producing governorates of the U.A.R., tests show importance in causing corn diseases, 53; survey of crown and root rots of alfalfa and red clover and the relative prevalence of associated fungi, 573; survival of selected fungi stored in liquid nitrogen for 17 months, 539
Fungicides: differential influence of variety on the pathogen-host-fungicide relationship is significant in formulating a control for Sclerotinia dollar spot of bentgrasses, 40; evaluation of, for control of crown rust, 130; fungistatic effect of 25 chemicals on the hyphal growth of cotton root rot fungus, 138; three highly effective for control of scab and other diseases of Pecan in southwestern Alabama, 185; addition of fungicides to infested soil in laboratory stimulated emergence of larvae from cysts of tobacco cyst nematode, 202; fungicidal and nutritional treatments for control of red thread disease of turfgrass in western Washington, 215; sprays for control of Fabraea leafspot of pear, 257; availability of "Fungicide-Nematocide Tests," Vol. 22, 319; for control of stripe smut of 'Merion' Kentucky bluegrass, 355; on highbush blueberry, for control of Glomerella cingulata, 401; effect of preplanting soil treatment on peanut losses and reduction of pod mycoflora, 461; granular formulations of soil fungicides evaluated for seedling disease

control in cotton, 487; chemical control of bacterial blights of dry field beans in Michigan by foliage sprays applied by ground and air equipment, 622; root application for control of downy mildew of opium poppy in India, 686; for control of peanut leaf rust and Cercospora leaf spots, several reduced incidence, sprays more effective than dusts in 1966 tests, 687; control of Botrytis neck rot in late harvested onions in New York by fungicidal dusts, 696; wax emulsion additives for control of storage decay in lemons, 781; systemic fungicides reasonably effective in preventing or suppressing infection of Cytospora on 'Redhaven' peach trees, 834; chemically induced changes in urediospores of Puccinia graminis tritici, 904; bulb treatment for the control of root-lesion nematode in Easter lily, 978; three low molecular weight carboxylic acids show no promise for prevention or therapy for Dutch elm disease, 1077

---: biphenyl: biphenyl-impregnated pads for storage decay control in Florida apparently not effective against one strain of citrus green mold, 700; new application method for control of citrus fruit rots, 58

---: captan dip treatment on crowns of crownvetch gave good control of storage decay, 820

---: 2,6-dichloro-4-nitroaniline (DCNA): gave excellent control of brown rot blossom blight of almonds, 578; effective in controlling sclerotial blight of sweetpotato, 191

---: 1,4-oxathiin derivative: for control of leaf and stem rust epidemics on wheat, 336; evaluation of two for control of seedling safflower rust, 586; two derivatives highly effective in reducing incidence of smutted kernels in pearl millet, 468

---: PCNB: in-furrow treatment with PCNB indicates some hope for control of corn head smut in Idaho, 626; to reduce losses from southern blight in peanut production, 441; to control southern blight on tomato and peanut, 314

---: phytoactin applications to western pine seedlings had no effect on normal mycorrhizae-root consociations, 1012

---: tetrachloroisophthalonitrile evaluated and compared with maneb for control of potato early blight under field conditions, 9

---: thiabendazole: tested in resin-solvent and wax-emulsion coatings for control of Penicillium digitatum in navel oranges, 51; as a postharvest fungicide for Florida citrus fruit, 95

---: several tested together: captan and maneb effective in control of leaf spot diseases of seedlings of shrub row plants skunkbush, sumac, and buffaloberry, 700; for control of Dutch elm disease, trials with captan and other fungicidal agents, 533; overtreatment of mercury-treated cottonseed with chloroneb increased plant populations, but not so effective as captan-PCNB, 558; chemotherapeutic effects of cycloheximide thiosemicarbazone, phytoactin, and 8-quinolinol benzoate with and without DMSO and pruning, on Cytospora canker disease of prune trees, 340; DL-β-methylaspartic acid (MAA) or DL-β-aminobutyric acid as side dressing on soil surface gave best control of Aphanomyces root rot of peas, 125; methyl bromide and chloropicrin effective in control of charcoal root disease in white fir and ponderosa pine nursery seedlings, 671; Semesan and Ceresan rhizome soak for reduction of Botrytis rhizome rot of Iris, 83; 45% thiram + 10% 2-chloro-4-(hydroxymercuri)phenol (T-OM) most effective turf fungicide tested for control of Fusarium blight of turfgrass, 839; ziram or mixture of nickel sulfate and maneb effectively controlled leaf blight of wheat in India, 950

Fusarium spp.: 11; 174; 376; 473; 601; 826

--- moniliforme: 540

--- nivale: 810; 815

--- oxysporum: 98; 629; 736

--- --- conglutinans race 2: 136

--- --- apii: 888

--- --- lycopersici: 645

--- --- perniciosum: 148

--- --- pisi: 888

--- roseum: 820; 839

--- solani: 531; 900

--- --- minus: new record on coffee in India, 758

--- wilt (see Wilt, Fusarium): Fusarium wilt infection of apparently healthy mimosa, 148

--- yellows: of radish, use of resistant varieties proved most effective means of controlling, 136

Fusicladium effusum: 185

Georgia: 461; 971; 1001

Geranium: bacterial leaf spot and stem rot in Pennsylvania, 61

Ghana: 941
Gloeosporium sp.: 212
Glomerella cingulata: 401
Gossypium spp.: 23 species and subspecies tested for any sources of resistance to several soilborne pathogens, 864
Grape: branch wilt, apparently a new disease found throughout grape-producing regions in Iraq, caused by Hendersonula toruloidea, 371; cane and leafspot phase of dead-arm disease found in Arkansas and West Virginia, 1038
Grasses: geographical distribution and prevalence of Helminthosporium victoriae, 110
---: bentgrass, Sclerotinia dollar spot, differential influence of variety on the pathogen-host-fungicide relationship, 40
---: Bermuda grass: pseudo-root-knot nematode on Tifgreen, first report in Alabama, 148; pseudo-root-knot nematode in Tennessee, 455; control of endo- and ecto-parasitic nematodes by hot water treatments, 905
---: Johnson grass: maize dwarf mosaic virus distribution in Iowa in 1966, 388; additional data on distribution and properties in Alabama, 385; plants and mosaic symptoms found in Louisiana, 108
---: Kentucky bluegrass: floral infections by Helminthosporium vagans, 752; role of nematodes in the "fading out" of bluegrass lawns, 457; 'Merion" Kentucky bluegrass, chemical control of stripe smut, 355
---: turfgrass: fungicidal and nutritional treatments for control of red thread disease of, 215; Fusarium blight of, and effectiveness of chemical control, 839; Bermuda turf, growth response and nematode reduction as influenced by nematocide applications, 562
---: zoysia: fungal diseases of zoysia in the Washington, D.C. area, 11; new severe blight disease developed under abnormal environmental conditions, caused by Fusarium spp., 376; rust, no known previous record in Louisiana, 726; rust of zoysia in Arkansas, first known report of, 893
Greece: 75; 936
Guar: leaf spot of, in Texas and Oklahoma, believed first report in United States, 182
Guatemala: 44; 265; 267
Gymnosporangium clavariiforme: 143

Hawaii: 1051
Helicobasidium compactum: 267
Helicotylenchus dihystera: 809
Helminthosporium avenae: 120
--- tetramera: 11
--- vagans: 752
--- victoriae: 470; geographical distribution and prevalence, 110
Hemlock (see Tsuga)
Hendersonula toruloidea: 371
Herbicides: unexpected preplant herbicide damage in sugarbeet possibly contributed to by plant pathogens, 86
Heterodera glycines: 771; 775; 823
--- tabacum: 609; tobacco cyst nematode, fungicides stimulated emergence of larvae from cysts, 202
--- trifolii: 1017
Hibiscus: foliar nematode Aphelenchoides fragariae found to be pathogenic to Hibiscus rosa-sinensis in greenhouse trial, 988
Hirschmanniella spp.: 960
Holly (see Ilex)
Hops: nematodes on, soil fumigation and irrigation for control of, 794
Host range: of three most prevalent parasitic nematodes associated with forest nursery seedlings in Quebec, 91; selective host range for certain mechanically transmitted stone fruit viruses, 786
Hosts: 84 species of plants found to be hosts to 1 or more of among 30 species of plant-parasitic nematodes tested in Nigeria, 33
Hot water treatments (see Temperature)
Hypsoperine sp.: 905
--- graminis: 148; 455

Idaho: 310; 591; 626
Ilex crenata: 'Helleri': found susceptible to five root-knot nematode species, 581; phosphorate nematocides gave excellent control of root-knot nematodes on dwarf Japanese holly, 1035
Illinois: 351; 511; 974
Index to genera of pathogens listed in the "Index of Plant Diseases in the United States"--4 parts, 235, 323, 419, 515
India: 138; 189; 480; 758; 764; 800; 950; 1079
Iowa: 388; 389; 438; 900; 1069
Iran: 595; 976
Iraq: 371; 690; 809
Iris: Botrytis rhizome rot, fungicide treatments for control of, 83; severe iris mosaic virus, symptomless occurrence in bulbous iris and its relation to color breaking in flowers, 952; yellow mosaic virus on, spread reduced by oil sprays in Israel, 318
Israel: 318; 373; 414; 733; 967

Juniperus communis: witches'-brooms caused by Gymnosporangium clavariiforme, taxonomic similarity of G. clavariiforme and G. gracile compared, 143

Kentucky: 731; 866
Key: of the most common species of the genus Alternaria, 295

Larch (see Larix)
Larix laricina: dwarf mistletoe on, in Minnesota, 418
Late blight (see Blight)
Leaf scald: of sugarcane, isolates thought to be Xanthomonas albilineans from sugarcane infected with leaf scald disease proved to be pathogenic, 491
Leaf spot: of annual phlox caused by Stemphylium botryosum, 741; on citrus, not previously reported on citrus in Florida caused by Cladosporium oxysporum, 1070; of french beans, caused by new virulent form of Phaeoisariopsis griseola in Tanzania, 276; of greenhouse plants due to soot damage, 712; of pansy, caused by Centrospora acerina reported in southern United States for first time, 971; of physiogenic origin on alfalfa and clover, 619; of shrub row plants skunkbush sumac, and buffaloberry, can be controlled by captan or maneb, 700
---: bacterial: angular, of cotton, absence of contamination in cotton gins 5 years after field eradication in San Joaquin Valley of California, 1066; of Guar, in Texas and Oklahoma, believed first report in United States, 182; holcus, on corn, epidemic in east-central Nebraska in 1966, 396; of strawberry in Kentucky, 353
---: Cercospora, on peanut, several fungicides reduced incidence of both leaf spots and rust in Texas, 687
---: Fabraea leafspot of pear, fungicidal control of, 257
Legumes: forage: resistance of several potential forage legumes to Meloidogyne species tested in Georgia, 1001; 76 species of Leguminosae tested for Heterodera trifolii cyst production, 1017
Lenzites trabea: 669
Lespedeza: 'Kobe' lespedeza unsuitable variety for differentiating isolates of soybean cyst nematode, 775
Lettuce: arabis mosaic virus in England, seed transmitted, 883; tomato spotted wilt virus occurring naturally in Argentina for first time, 225
Lily: Easter: bulb treatments for the control of root-lesion nematode Pratylenchus penetrans, 978; after soil fumigation with nematocides, lily bulbs showed significant increase in size and weight, 549
Liriodendron tulipifera: Cylindrocladium scoparium in a 27-year-old plantation, first report of extensive root damage and mortality caused by this fungus in trees beyond the seedling stage, 38
Longevity: survival of selected fungi stored in liquid nitrogen for 17 months, 539; overtreatment of mercury-treated cottonseed with chloroneb did not affect germination for periods up to 7 months, 558; attempt to extend the longevity of burrowing nematode, Radopholus similis, with citrus root exudate, 675; effect of sealed and unsealed containers on longevity of dried cultures of microorganisms, 747; for maximum longevity of fungi, actinomycetes and bacteria in storage, dry at low temperature and store in sealed containers in refrigerator at 10 to 13°C, 1055
Longidorus spp.: 30
--- elongatus: 1028
--- sylphus: 809
Loose smut: of oats, distribution and prevalence of virulence for Victoria oats, 846; pathogenicity in races of Ustilago avenae for adaptation to changing host varieties of oats, 844
Louisiana: 67; 108; 167; 726; 823

Macrophomina phaseoli: 671
Maine: 608
Malaysia: 443
Mango: Diplodia stem-end rot on, mode of infection and prevention, 744
Manitoba: 899
Maryland: 345
Meloidodera sp.: 264
Meloidogyne spp.: 373; 482; 581; 614; 1001; 1035
--- hapla: 103; 164; 736
--- incognita: 219
--- --- acrita: 29; 470; 720
--- javanica: 861
--- naasi: 974
Methods and techniques: an agar-layer method useful in detecting anitbiotic-producing microorganisms against Pythium graminicola, 333; use of aluminum foil as protective device to control spread of barley yellow dwarf virus in winter oats not effective, 354;

use of sheep to produce large quantities of high titer antiserum against plant pathogenic bacteria, 50; modified Baermann funnel technique described as good method for obtaining aseptic inoculum of nematodes, Aphelenchus avenae, 367; new method of applying biphenyl for control of fruit rots of citrus, 58; constant temperature bath for pot-grown plants, 171; effect of sealed and unsealed containers on longevity of dried cultures of microorganisms, 747; quick destaining procedure for showing contrast between nematodes and root tissue, 511; hill plots useful as relatively economical and reliable field-testing method for use with promising chemicals against crown rust of oats, 130; incubation of hydrogen peroxide-treated root samples in plastic bags to increase nematode yields from citrus roots, 1024; method for obtaining quick infection with Sclerospora species, 1009; three investigative methods for predicting late blight of potato based on rainfall, temperature and relative humidity in Greece, 936; comparison of three techniques for extraction of needle nematode from soil, 1028; storage of selected fungi in liquid nitrogen, 539; early cereal aphid and barley yellow dwarf detection in Iowa by utilizing documented periods of low-level jet winds, 527; mist culture, seems well suited for investigation of various root phenomena and root development, 1074; a soil-temperature control system employing air as the heat-transfer medium, 873; a simple field method for inoculation of white pine blister rust cankers with purple mold spores to suppress rust fungus, 89; technique for labeling Ceratocystis ulmi with radiocarbon, 803; seed growth pouches used for root-knot nematode resistance tests, 482

Michigan: 261; 476; 622

Microxyphiella sp.: on cotton, no known previous record, 743

Millet: first record of Piricularia setariae as a seedborne pathogen in Setaria italica, 138; two 1,4-oxathiin fungicides highly effective in reducing incidence of smutted kernels in pearl millet, 468

Mimosa: Fusarium wilt of wilt-resistant cultivars, 148

Minnesota: 418; 679

Mississippi: 30; 1044

Missouri: 377; 760

Monilinia fructicola: 750

--- laxa: 578

Montana: 702; 791

Mulberry: Fusarium canker on paper mulberry, 531

Nebraska: 396

Nectria galligena: first report of the Nectria fruit rot phase in European canker orchards in California, 278

"Needle curl": of loblolly pine seedlings, associated with excessively high temperatures, 805

Needle fungus: first report on Engelmann spruce in Montana, 791

Nematocides: availability of "Fungicide-Nematocide Tests," Vol. 22, 319; effect of variety and fumigation with methyl bromide on nematode populations in wheat, oats, and rye in Florida, 915; phosphorate nematocides for control of root-knot nematodes on dwarf Japanese holly and Japanese boxwood very effective, 1035

---: DBCP: DBCP and 1,3-D: tested for control of nematodes on hops, 794

---: DD: reduction of nematode population and increased tree growth in response to treatment of planting site of cherry orchard with dichloropropene-dichloropropane, 398; gave good control of plant-parasitic nematodes in corn, however, did not result in significant increases in yields, 15

---: several tested together: cynem and dimethoate gave best control of stem nematode Ditylenchus dipsaci in alfalfa seedlings, 973; dips, for control of nematodes on bare-root plants, 103; influence of applications on nematode reduction and upon growth response of Bermuda turf, 562; on Kentucky bluegrass, partially controlled stylet-bearing nematodes, 457; effective control of nematodes on citrus seedlings using chemical bare-root dips, 995; effect of five nematocidal treatments for control of root-lesion nematode and upon growth of bulbs, 549; in combination with fungicidal emergence stimulants, nematocides may help reduce population of tobacco cyst nematodes in field, 609; for control of root knot of tomatoes, 219; fumigation studies for control of nematodes associated with roots of wheat in Nebraska, 637

Nematodes: associated with azalea in Florida, first report of 14 genera and 20

species associated with azalea, 46; on Bermuda turf, population reduction as influenced by nematocide applications, 562; soil preplant fumigation resulted in greater growth of cherry trees in treated soil, 398; quick destaining procedure for showing contrast between nematodes and root tissue, 511; lily bulbs greater in size and weight after soil fumigation with nematocides, nematode population also greater, 549; oat varieties showed stability of reaction to Helminthosporium victoriae and Meloidogyne incognita acrita in dual inoculations, 470; first report in Iraq of four nematode species being associated with diseased condition of sugarcane, 809; population dynamics and fumigation studies of certain nematodes associated with roots of wheat in Nebraska, 637; effect of variety and fumigation on nematode populations in oats, wheat, and rye in Florida, 915; on hops, soil fumigation by injection and irrigation for control, 794; host range and reproduction of three most prevalent plant-parasitic nematodes associated with forest nursery seedlings in Quebec, 91

---: plant-parasitic: controlled by soil fumigation with D-D, however, corn yields did not increase significantly, 15; of rice, constitute economic problem in Louisiana rice fields, 167; 84 species of plants found to be hosts to 1 or more of among 30 species of nematodes tested in Nigeria, 33; survey for presence of plant-parasitic nematodes on stone-fruit rootstocks from an experimental orchard, in Israel, 373

---: Aphelenchus avenae failed to parasitize roots of seven species of pine seedlings under non-aseptic or aseptic conditions, 367

---: burrowing: annual cycles of increase and decrease in populations of Radopholus similis occur in Florida citrus groves, 1031; attempt to extend the longevity of burrowing nematode, Radopholus similis, with citrus root exudate unsuccessful, 675; effect of temperature and hydrogen peroxide on extraction of nematodes from citrus roots, 1024

---: citrus: distribution in Florida, 317; effective control on infected seedlings by using nematocidal bare-root dips, 995

---: cyst: cyst production of Heterodera trifolii on some legumes, 1017

---: dagger: transmission of peach yellow bud mosaic virus by Xiphinema americanum to peach, plum, and apricot seedlings, 841

---: ecto- and endo-parasitic: influence of temperature, photoperiod, and nutrition on reproduction, male-female-juvenile ratio, and root to soil migration of Pratylenchus penetrans, 78; control of ecto- and endoparasitic nematodes by hot water treatments, 905

---: foliar: Aphelenchoides fragariae found pathogenic to Hibiscus rosa-sinensis, 988; two new fern hosts of, in Florida, 508

---: Meloidodera sp., a new nematode found infecting mycorrhizae of Douglas-fir, 264

---: needle: some factors influencing extraction of needle nematode from soil, 1028; 2 species, Longidorus brevicaudatus and a new Longidorus sp., found in Sudan for first time, 30

---: Pratylenchus brachyurus, pathogenicity to citrus established, 569

---: pseudo-root-knot: of Bermuda grass in Tennessee, 455; on Tifgreen Bermuda grass, in Alabama, new distribution, 148

---: root-lesion: first record of nematode infestation of eucalyptus trees in S. America, 791; bulb treatment for control of root-lesion nematode Pratylenchus penetrans in Easter lily, 978

---: reniform: found in five southeastern States reported for first time in South Carolina, 557

---: rice root: study of in Thailand, 960

---: root-knot: amendments of sawdust + NPK fertilizer to soil suppressed infestation and increased yield of okra and tomatoes, 861; of barley, first report of occurrence in Illinois, 974; of cotton, associated with damage to grain sorghum in west Texas, 29; pathogenicity of Meloidogyne incognita acrita to Cucumis species in greenhouse screening trials, 720; elimination of by chemical bare-root dips or soil drenches, 103; first report of Meloidogyne hapla on Euphorbia lagascae, 164; susceptibility of potential forage legumes to Meloidogyne species, 1001; susceptibility of Ilex crenata 'Helleri' to five root-knot nematode species, Meloidogyne spp., 581; phosphorate nematocides as soil drenches for dwarf Japanese holly and Japanese boxwood gave excellent

control of, 1035; on Pinus elliotii seedlings in Florida, 455; use of seed growth pouches for resistance tests, 482; survey for resistance of stone fruit rootstocks from an experimental orchard in Israel, 373; on tomato, nematocides superior to crop residues for control of, 219; additional Vicia species found resistant to root-knot nematodes, 614
---: saprozoic: laboratory studies suggest possibility of significant role in disseminating or transmitting bacteria and fungus spores, 98
---: soybean cyst: first reported occurrence in Louisiana, 823; nature of soybean resistance to, 771; 'Kobe' lespedeza unsuitable for reproduction of isolates of Heterodera glycines, 775
---: stem: in alfalfa, an evaluation of several techniques for screening alfalfa for resistance to stem nematodes, 651; good control of Ditylenchus dipsaci in alfalfa seedlings with cynem and dimethoate, 973
---: stylet bearing: in Kentucky bluegrass, stylet-bearing nematodes appear to contribute to "fading out" of bluegrass, 457
---: tobacco cyst: effect of fungicides on hatching, survival, and nematocidal killing of eggs of Heterodera tabacum in the field, 609; stimulation of emergence of larvae from cysts of tobacco cyst nematode by adding fungicides to infested soil, 202
New diseases: Cylindrocladium scoparium in a 27-year-old yellow-poplar plantation, first report of extensive root damage and mortality caused by this fungus in trees beyond the seedling stage, 38; a graft-transmissible "red-ring" disease described, possibly a new disease of apple, 281; severe blight disease of zoysia developed under abnormal environmental conditions, caused by Fusarium spp., 376; branch wilt of grape in Iraq, caused by Hendersonula toruloidea, 371
---: in India: Rhizopus rot of chicory, 189; twig blight of Vinca rosea, 480
New distribution: list of fungus pathogens found on cultivated plants and weeds in Chile, not previously recorded on these plants, 212; plant diseases previously unreported from Hawaii, 1051; occurrence of a yellows-type disease in periwinkles in North Africa, 416
---: bacterial canker of tomato in Chile, 396
---: Cladosporium leaf spot, not previously described on citrus in Florida, 1070
---: collar canker of coffee trees, spread in Guatemala, 267
---: Dutch elm disease found in southern Arkansas represents one of the more southern points of its distribution nationally, 71
---: fire blight disease, first authentic report of, on continental Europe, 3
---: maize dwarf mosaic virus found in 4 new locations in Iowa in 1966, 388
---: nematodes: Longidorus brevicaudatus and a new Longidorus sp. found in Sudan for first time, found on Jew's mallow and spearmint, 30; pseudo-root-knot nematode in Alabama, 148; of Bermuda grass in Tennessee, 455; first reported occurrence of soybean cyst nematode in Louisiana, 823
---: pansy leafspot reported in southern United States for first time from Georgia, 971
---: Phacidiopycnis pseudotsugae, first report of affecting hemlock naturally, 807
---: Piricularia setariae, first record as a seedborne pathogen in Setaria italica from India, 138
---: scab of celeriac in the United States (Wisconsin) for first time, 427
---: Scleroderris pine canker, first record on seedling stock in North American forest nurseries, 357
---: tomato spotted wilt virus occurring naturally in lettuce in Argentina, 225
---: Tranzschelia discolor and Stereum purpureum found on peach and apricot trees in Chile, previously unreported, 212
New hosts: in Chile, additional records of fungi not previously reported on particular host plants, 213; 14 genera and 20 species of plant-parasitic nematodes associated with azalea for first time, 46; plant diseases previously unreported from Hawaii, 1051;
---: Armillaria root rot not previously reported from California, 390
---: Botryosphaeria canker on Johathan apple trees in Washington, 593
---: poplar spotted anthracnose on Populus angustifolia, found in Oklahoma, 785
---: Microxyphiella sp., first report of, on cotton, 743
---: nematodes: Meloidogyne hapla on Euphorbia lagascae, 164; root-knot nematode, found on Pinus elliottii, 455; first report of root-lesion

nematode on eucalyptus, 791
---: Stemphylium botryosum, on Phlox drummondii in New Jersey nurseries, 741
---: Verticillium nigrescens, on cotton, 760
---: Viola adunca and Agropyron smithii non-rosaceous hosts of Phragmidium (Uredinales), 766
New Jersey: 349; 709; 850; 964
New York: 195
New Zealand: 727
Nigeria: 33; 174
Nitrogen: liquid, survival of selected fungi stored in liquid nitrogen for 17 months, 539
North Africa: 416
North Carolina: 409; 431
Nutrients: effect of low calcium levels of nutrition upon susceptibility of tobacco seedlings to black shank disease, 641; addition of hydrated lime to soil inhibited development of Fusarium wilt (race 2) in tomato (variety Manapal), 645; minimum level of nitrogen necessary for good turf growth helps reduce incidence of red thread disease in western Washington, 215; influence on reproduction, male-female-juvenile ratio, and root to soil migration of Pratylenchus penetrans, 78

Oak: decay in hardwood trees often inhibited or halted after wounds heal, 600; vascular mycosis of, in Russia, 739
---: wilt: antibiotic treatments for possible control of, 160; evaluation of oak bark beetle as a vector, 755; wilt mats and pressure cushions found on white oak for first time, West Virginia, 430
Oats: nematode populations in Florida reduced in fumigated fields, 915; stand loss of small grains in Florida, 919; stability of reaction of oat varieties to Helminthosporium victoriae and Meloidogyne incognita acrita in dual inoculations, 470; wild oat species show promise as sources of disease resistance for improvement of cultivated oats, 391
---: crown rust: use of hill plots in evaluation of fungicides for control of, 130; most prevalent races of crown rust in Texas for 1952-1965 tabulated, relating incidence to estimated acreages of oat varieties grown, 133; a quantitative test of tolerance of experimental lines of oats to a mixture of common races of crown rust, 947; selections of Avena sterilis tested at four locations showed resistance to race 264, 120
---: loose smut: distribution and prevalence of virulence for Victoria oats, 846; race population dynamics in Ustilago avenae as related to persistence of loose smut in oats, 844
---: powdery mildew, observations of differences in resistance and susceptibility of newer winter varieties under epidemic conditions in the field, 30
---: stem rust, new and highly virulent race of oat stem rust in Israel designated race 41, 967
---: virus: barley yellow dwarf virus, 'Blount' and several Tennessee breeding lines show some tolerance to, 908; soilborne mosaic virus, resistance observed in 43 selections of Avena sterilis tested at Quincy, Florida, 120
Oil: spray, for reducing yellow mosaic virus in Irises in Israel, 318; reduction of lesion nematodes in plant-derived oil-amended soils, 1021
Oklahoma: 785
Okra: amendments of sawdust and balanced NPK fertilizer to soil suppressed root-knot infestation and increased yield in India, 861
Onion: asparagus rust: all commercial onion varieties grown in southwestern Idaho appear susceptible to, 542; Botrytis neck rot controlled in freshly harvested onions by fungicidal dusts in New York, 696
Ontario: 357
Ophiostoma (Ceratocystis) roboris: 739
Oregon: 264; 281
Ozone: Mast ozone analyzers and ozone-sensitive tobacco plants used to detect presence of oxidants on the Delmarva Peninsula, 493; possible cause of tip burn of carnation, 793; appears to be principal air pollutant in New Jersey, 850; related to occurrence of weather fleck on Florida cigar-wrapper tobacco in 1966, 72

Paint: fungicidal, for aiding prevention of Diplodia stem-end rot of mango, 744
Pakistan: 743
Pansy: pansy leafspot reported in southern United States for first time, 971
Pathogenicity: of Phacidium abietis on several conifer species, 678; studies with isolates from sugarcane infected with leaf scald disease suspected to be Xanthomonas albilineans, proved to be pathogenic, 491
Pathogens: associated with root rots of conifers in South Australian

nurseries, 473; index to genera of pathogens listed in the "Index of Plant Diseases in the United States"--4 parts, 235, 323, 419, 515
Pea: Aphanomyces root rot, comparison of chemical soil treatments and soil amendments for control of, 125; Fusarium wilt of celery, caused wilt of garden pea, 888
Peach (see also Prunus): bacterial canker transmitted by infected budwood, 666; use of systemic fungicides for prevention and cure of Cytospora canker disease, 834
---: bacterial spot: control with Terramycin, 345; superior control by antibiotics Terramycin and streptomycin, 349; effect of Terramycin and dodine-captan mixture on prevention of peach bacterial spot, 351
---: brown rot, result of direct penetration by germinating conidia of, 750
---: rust, caused by Tranzschelia discolor in Chile for first time in 1966, 212
---: silverleaf, caused by Stereum purpureum, first report in Chile, 212
---: viruses: incidence in Washington peach orchards of latent viruses, 1063; maintaining virus-free cultivars of peach in Michigan, 261; virus resembling peach yellows found in Israel, 733
Peanut: chlorosis due to iron deficiency, 464; pod breakdown, relation to several pathogenic organisms, and the competition of Trichoderma viride to breakdown, 601; drying of topsoil for day or two between irrigations appears to impede Pythium and combat pod rot in Israel, 414; effect of preplanting soil treatments on peanut losses and reduction of pod mycoflora, 461
---: peanut leaf rust and Cercospora leaf spot, fungicides reduced incidence, sprays more effective than dusts in 1966 tests, 687
---: Cylindrocladium rot, plant age and soil temperature interact to affect disease severity, 986
---: root rot, varietal response to Sclerotium bataticola in India, 649
--- southern blight: effective control of Sclerotium rolfsii by deep plowing and PCNB treatment on irrigated fields in Texas, 314; yields of peanuts significantly increased by planting in medium or high beds and application of PCNB to reduce losses from southern blight, in Texas, 441
---: stunt virus: drastically reduced yield and quality of peanuts in commercial fields in Virginia, 856; stunt disease survey in Virginia in 1966, 169
Pear: Fabraea leafspot of, suggested resistant pear variety plus limited spray schedule for effective and economical control of, 257; fire blight, first authentic report of disease on continental Europe, 3; mechanical transmission of pear viruses from pear to Chenopodium amaranticolor, 283
Pecan: scab control and other diseases, three fungicides highly effective for, 185
Pellicularia filamentosa: 67
--- sasakii: 682
Penicillium spp.: 461
--- digitatum: 51; 95; 700; 781
--- italicum: 781
--- rubrum: 669
Pennsylvania: 61; 941
Pepper: bacterial spot, effect of growth-regulating and other compounds on, 320; 497
Peppermint: Verticillium wilt of, effectively reduced in muck soils after 5 years' continuous cropping with corn or after reed canary grass sod, 449
Peridermium harknessii: a white spore race, 306
Periwinkle (see Vinca)
Peronospora arborescens: 686
Petunia: transmission of exocortis virus of citrus, 868
Phacidiopycnis pseudotsugae: 807
Phacidium abietis: 678
Phaeoisariopsis griseola: 276
Philippines: 692: 877
Phlox: leaf spot, caused by Stemphylium botryosum, 741; perennial, root, crown, and basal stem rot caused by Thielaviopsis basicola and soil temperature, moisture, and pH, 436; studies of mosaic virus on, 768
Phoma apiicola: 427
--- lingam: 566
Phomopsis citri: 95
--- viticola: 1038
Photoperiod: influence on reproduction, male-female-juvenile ratio, and root to soil migration of Pratylenchus penetrans, 78; 16-hour day showed significant increase in quantity of sporangia of Phytophthora phaseoli produced on 'Fordhook' lima bean stems over a 12-hour day, 253
Phragmidium sp.: 766
Phymatotrichum omnivorum: 138; 331; 926
Physalis minima: ground cherry, as suspect of bacterial spot, Xanthomonas

vesicatoria, 659
Physalospora obtusa: 230
--- rhodina (Diplodia natalensis): 744
Physiological diseases: chilling-injury syndrome in potato tubers, 1014; stylar-end rot of limes, reduced by use of organic sprays, 1071
Phytobia pruni: 941
Phytopathological Monographs available: 322
Phytophthora spp.: 929
--- cactorum: 212
--- cinnamomi: 998
--- infestans: 75; 936
--- (?) meadii: 443
--- megasperma: 679
--- palmivora: 443
--- parasitica nicotianae: 641; 669; 731
--- phaseoli: 253
Picea spp.: three most prevalent plant-parasitic nematodes associated with nursery stock in Quebec reproduced better on spruces than on pines, 91; Englemann spruce, first report of needle fungus (Bifusella crepidiformis) in Montana, 791
Pinus spp.: cone rust, rapid spread of this main problem of pine forests in Guatemala, 44; under non-aseptic and aseptic conditions the nematode Aphelenchus avenae did not parasitize roots of seven species of pine seedlings, 367; occurrence and pathogenicity of Scleroderris lagerbergii in Quebec, 584; needle blight (Dothistroma pini), distribution and significance in New Zealand, 727; western white pine, applications of phytoactin to pine seedlings had no effect on normal mycorrhizae-root consociations, 1012
--- banksiana (jack pine): first authentic record of infection by eastern dwarf mistletoe in Manitoba, 899
--- elliottii (slash pine): a new host for root-knot nematode Meloidogyne javanica, 455
--- lambertiana (sugar pine): simple field method for inoculation of blister rust cankers with purple mold spores to suppress rust fungus, 89
--- nigra var. austriaca (Austrian pine): extensive needle damage in New Jersey after period of high air pollution, 964
--- ponderosa (western yellow pine): effective control of charcoal root disease of nursery seedlings by fall soil fumigation, 671; dwarf mistletoe infected, distribution on South Rim of Grand Canyon, Arizona, 1049; western gall on, a white spore race, 306
--- radiata (Monterey pine): susceptibility of two seedlings to infection from contact with bishop pine mistletoe seeds in experimental testing, 856
--- resinosa (red pine): Scleroderris canker in New York plantations, 108; Scleroderris pine canker, first confirmed report on seedling stock, in Ontario, 357
--- strobus (eastern white pine): rate of decay from artificial inoculation with Fomes pini extremely variable, 829
--- taeda (loblolly pine): brown spot, field infection of pine seedlings with naturally produced inoculum of Scirrhia acicola, 552; comandra rust in eastern Tennessee, 4; antibiotic treatments for fusiform rust of, 223; fusiform rust galls on, effectiveness of surgical and antibiotic treatments upon sporulation, 505; "needle curl" associated with excessively high temperatures, 805
Piricularia oryzae: 243
--- setaria: 138
Plasmodiophora brassicae: 991
Pleurotus ostreatus: 600
Plum (see also Prunus): 'Shiro' plum a rapid greenhouse indicator for peach wart virus, 831
Poland: 3; 115; 117
Poplar (see Populus)
Poplar, yellow- (see Liriodendron)
Poppy, opium: partial control of downy mildew by root applications of certain fungicides, 686
Populus: canker disease, first report of occurrence in Iowa, 900
Potato: distribution of Verticillium spp. in commercial potato-growing regions of Maine, 608; early blight, evaluation of tetrachloroisophthalonitrile for control, 9; physiological breakdown in potato tubers stored at 32°F in air occurred as chilling-injury syndrome, 1014; purple-top (aster yellows virus) in Guatemala, 265; "running out" in noncertified whole potatoes used for seed due to seed rather than cultural practices, 298
--- late blight: 11 years of data on weather and occurrence of disease analyzed for value in forecasting, Thessaloniki, Greece, 75; relation between weather data and blight in forecasting occurrence in Greece, 936
Powdery mildew: on Brassica campestris var. sarson and Brassica juncea in India, first record of perithecial stage of the fungus, 800; of cucurbits, identity of powdery mildews of cucurbits in India, 1079; of winter oats,

evaluation of differences in resistance and susceptibility of newer varieties under epidemic conditions in the field, 30; of wheat, fluorochemical liquid FC-43 found effective as carrier for conidia of Erysiphe graminis to plant surfaces, 447; of wheat, 1966 epiphytotic in eastern Washington, 94
Pratylenchus brachyurus: 569; first report of root-lesion nematode on eucalyptus, 791
--- penetrans: 15; 78; 978; 1021
--- zeae: 809
Preservation: of Xanthomonas albilineans in distilled water, 739
Pruning: tools: sodium hypochlorite as disinfectant for, in control of fire blight, 753
Prunus (see also common names): new records of fungus diseases of fruit crops in Chile, 212; survey of stone fruit rootstocks from an experimental orchard for presence of plant-parasitic nematodes and resistance to root-knot nematodes in Israel, 373; excellent control of brown rot blossom blight on almonds obtained in California from DCNA spray, 578
---: canker: chemotherapeutic effect of cycloheximide thiosemicarbazone, phytoactin, and 8-quinolinol benzoate with and without DMSO and pruning, on the Ctyospora canker disease of prune trees, 340; evaluation of first and second year effects of cycloheximide thiosemicarbazone treatments on Cytospora canker disease of 'Italian' prune trees, 655; prevention and cure of Cytospora canker disease with systemic chemicals, 834
---: viruses: a selective host range for, 786; apple chlorotic leaf spot and tobacco mosaic viruses in cherry, 823; cross-transmission studies with Citrus and Prunus viruses, 64; Lambert mottle (virus), effect of temperature on symptom development of, 269; peach wart virus detection possible on 'Shiro' plum in 4 weeks, 831; peach wart virus symptoms in several Prunus spp., and peach and nectarine varieties, 42; transmission of peach yellow bud mosaic virus to peach, apricot, and plum seedlings by dagger nematode, 841
Pseudomonas lapsa: 764
--- phaseolicola: 20; 310; 544; 591; 622; 661
--- syringae: 182; 396; 666
Pseudopityophthorus spp.: 755
Pseudotsuga menziesii: a new nematode Meloidodera sp. found infecting mycorrhizae of Douglas-fir, 264; occurrence of needle blight in Chile, 454
Psyllids: citrus leaf-mottle-yellows disease in Philippines transmitted by psyllid, Diaphorina citri, 692
Puccinia arachidis: 687
--- asparagi: 542
--- carthami: 586; 589
--- coronata avenae: 120; 130; 391; 947; tabulation of most prevalent races of crown rust in Texas for 1952-1965, 133
--- graminis: 358; 362; 447
--- --- avenae: 391
--- --- tritici: 248; 536; 904
--- recondita: 358; 362
--- striiformis: 362; 702; 911; 1041
--- zoysiae: 726; 893
Pumpkin (see Cucurbits)
Pythium spp.: 414; 929
--- aphanidermatum: 919
--- graminicola: 333
--- myriotylum: 601
--- ultimum: 864

Quarantine: virus detection in vegetatively propagated plant introductions under quarantine in the United States, 1957-1967, 715
Quebec: 91; 573; 584; 678

Races: a white spore race of the western gall rust fungus on ponderosa pine, 306; interaction of race I and race II of Colletotrichum orbiculare on watermelon, 885; Race 2 of Fusarium oxysporum lycopersici, liming of soil reduced incidence of wilt in tomato (variety Manapal), 645; distribution of race 1 of Phytophthora parasitica nicotianae in Kentucky and Tennessee, 731; screening tests of crucifers for resistance to race 6A of Plasmodiophora brassicae, 991; of Pseudomonas phaseolicola, several bean varieties tested for reaction to, 20; wheat bunt, a new race of Tilletia caries (designated X-1) with pathogenic range that suggests that it carries at least one gene for virulence that has not been observed in any other race of common bunt, 287
---: rust: new and highly virulent race of oat stem rust in Israel, 967; prevalence of Puccinia coronata avenae races in relation to estimated acreages of oat varieties in Texas, 1952-1965, 133; of Puccinia graminis tritici identified from Texas collections, 1926-1965, 536; physiological races of Puccinia recondita and P.

graminis in Rumania in 1964 and 1965, 358
Radiocarbon: technique for successfully labeling conidia of Dutch elm disease fungus, 803
Radopholus similis: 995; 1024; 1031
Red stalk rot (see Rot)
Red thread: of turfgrass in western Washington, fungicidal control of, 215
Relative humidity: RH of 0 to 5% most suitable to maintain viability of dried cultures of microorganisms, 1055
Resistance: sources of resistance in the genus Gossypium to several soilborne pathogens, 864; of mimosa to Fusarium wilt may be reduced by age, cold, or other factors, 148; of new radish (White Spike) to Fusarium yellows, 136
Rhizoctonia solani: 11; 86; 158; 212; 225; 601; 864; 872
Rhizopus stolonifer: 189
Rhode Island: 1053
Rice: bacterial leaf streak, in India, 928; blast, spread in small fields, 243; hot water seed treatment for eradication of inoculum of Xanthomonas oryzae, 882; plant-parasitic nematodes of, constitute economic problem in Louisiana rice fields, 167; studies of rice-root nematodes in Thailand, 960; sheath blight, in Taiwan, 20 strains of Pellicularia sasakii detected and studied for first time, 682
---: virus: yellow dwarf virus, preliminary note on occurrence and transmission in India, 1040; maize dwarf mosaic virus inoculation caused only localized infection, and exposure to MDMV in field did not cause infection, 381; rice tungro virus, transmission studies of the two strains, 877; virus-like disorder of rice, leaf yellowing in India, positive transmission by Nephotettix bipunctatus, 300
Root: exudate of citrus, of no significance for extending longevity of burrowing nematode in absence of contact with living roots, 675
Rots: fruit, of citrus, new method of applying biphenyl for control, 58; leaf and pod of rubber tree by two Phytophthora spp., confirmed for first time in Malaysia 1966, 443
---: bacterial soft rot, commercial practice of field washing mature-green tomatoes increased incidence of bacterial soft rot, 151
---: bacterial stem, of geraniums in Pennsylvania, 61
---: cotton boll: Pellicularia filamentosa found to be a primary cause, 67; no loss of viability or virulence after storage of boll rot organisms on infected cotton bolls for 8 years, 855
---: Botrytis rhizome, on Iris, 83
---: brown stem, of soybeans, incidence in Iowa, 438
---: red stalk, of sorghum, root injury appears to hasten death of stalk cells which increases susceptibility of plant to red stalk rot, 957
---: Rhizopus, new disease of chicory in India, 189
---: root: method of detecting antibiotic-producing colonies of fungi against root rot fungus Pythium graminicola, 333; of snapbean, effect of planting depth and asphalt mulch upon incidence of Rhizoctonia root rot, 158; Phytophthora, of blueberry, relative susceptibility of blueberry cultivars to, 998; of red and white clover, effect of cutting height upon incidence of root rot, 826; of conifers in South Australian nurseries, organisms associated with, 473; and stalk, of corn, D-D not recommended for control, 15; of cotton caused by Phymatotrichum omnivorum, fumigants for control of, 331; of cotton, fungistatic effect of chemicals in inhibiting growth of cotton root rot fungus may be applicable to other soil fungi, 138; of pea, best control obtained with DL-β-methylaspartic acid (MAA) or DL-β-aminobutyric acid as side dressing on soil surface for Aphanomyces root rot of peas, 125; of peanut in India, evaluation of varietal response, 649; root, crown, and basal stem rot of perennial phlox caused by Thielaviopsis basicola and soil temperature, moisture, and pH, 436; root and stem, of soybean, caused by Rhizoctonia solani, 872; brown root rot of tobacco, study gives supporting evidence that Rhizoctonia solani is strongly implicated in brown root rot of tobacco, 225; tests of pathogenicity on important shade and timber trees in the South, 500; root and crown, of zoysia, fungi associated with, 11
---: soil, of sweetpotato, development in Olivier silt loam as affected by lime or sulfur applications, 271
---: stalk, of corn, influence of cultural practices on the incidence of, 540
---: stylar-end rot of limes, use of organic spray materials reduced incidence, 1071

Rotylenchulus reniformis: 557
Rubbertree: outbreak of leaf fall and pod rot associated with both Phytophthora palmivora and P. (?) meadii confirmed for first time in Malaysia 1966, 443
"Running out": of potatoes, yield from noncertified whole potatoes used for seed significantly less than from cut certified seed potatoes, 298
Russia: 739
Russet: stem cavity, of apples, 302
Rust: usefulness of fluorochemical liquid FC-43 as a carrier for uredospores of Puccinia graminis evaluated, 447
---: asparagus rust, on fence row asparagus in southwestern Idaho, appears to be pathogenic to commercial onion varieties, 542
---: blister, of sugar pine, blister rust cankers inoculated with purple mold spores were markedly suppressed but seldom if ever completely controlled, 89
---: broadbean, with chocolate spot and bean yellow mosaic virus reached epidemic proportions in Iran 1966, 595
---: comandra, on loblolly pine in eastern Tennessee, recently found in widely separated plantations, 4
---: cone, of pine trees, rapid spread of this main problem of pine forests in Guatemala, 44
---: crown: of oats: resistance to virulent race 264 of crown rust and soilborne mosaic virus observed among 101 selections of Avena sterilis, 120; use of hill plots in evaluation of fungicides for control of, 130; tabulation of most prevalent races of crown rust in Texas, 1952-1965, relating their incidence to estimated acreages of oat varieties grown, 133; wild oat species showed resistance to race 264 of crown rust, 391
---: fusiform: of loblolly pines, antibiotic treatment of, 223; rust galls on loblolly pine, effectiveness of surgical and antibiotic treatments upon sporulation, 505
---: western gall, on ponderosa pine, a white spore race, 306
---: leaf: of peanut, several fungicides reduced incidence of both rust and leaf spots in Texas, 687
---: --- and stem: of wheat: physiological races of Puccinia recondita and P. graminis in Rumania, 358; control by 1,4-oxathiin derivative, 336
---: safflower: 21 collections of noncultivated species evaluated for resistance to rust and Verticillium wilt, 589; seed treatment with 1,4-oxathiin fungicides effectively controlled seedborne rust spores and reduced rust infection, 586
---: stem: of oats: wild oat species showed resistance to sulture of race 2AF of stem rust, 391; a new and highly virulent race of oat stem rust, designated race 41 in Israel, 967
---: ---: of wheat: chemically induced changes in urediospores of Puccinia graminis tritici, 904; races of Puccinia graminis tritici identified from Texas collections, 1926-1965, 536; some uredospore characteristics of importance for adequate prediction of field infection in experimental initiation of rust epidemics, 248
---: stripe: stripe, leaf and stem: of wheat, reactions of adult wheats to, in Rumania in 1965 as well as reaction of seedlings of six winter wheats from Rumania to stem rust cultures in United States in 1966, 362; comparison of root systems produced by healthy and stripe rust-inoculated wheat in mist-, water-, and sand-culture, 1074; induction of infection during summer in southeastern Washington, 911; single gene control of stripe rust resistance in wheat, 1041; effect on yield and its components of six varieties, 702
---: Tranzschelia discolor on peach, plum, and apricot in Chile for first time, 1966, 212
---: witches'-brooms as well as swollen stems on Juniperus communis are caused by Gymnosporangium clavariiforme, 143
---: of Zoysia: in Arkansas, first known report of, 893; first report in Louisiana, 726
Rye: anthracnose, in Florida, major disease limiting seed production, 923
---: stand loss of small grains in Florida, 919
---: effect of variety and fumigation on nematode populations in Florida, 915

Safflower: 21 collections of safflower species evaluated for resistance to rust and Verticillium wilt, 589; seedling rust, seed treatment with 1,4-oxathiin fungicides effectively controlled seedborne rust spores and reduced rust infection, 586
Scab: of celeriac (root celery) reported in U.S. (Wisconsin) for first time, 427; of cucurbit, reduction of by a 1-year

rotation with corn, 495; Fusarium, of sorghum in Nigeria, 174; of pecan, control of, in southwestern Alabama, 185
Scirrhia acicola: 552
Scleroderris lagerbergii: 108; 357; 584
Sclerophthora sp.: 1010
Sclerospora spp.: 1009
--- sacchari: 1010
Sclerotinia homoeocarpa: 40
Sclerotium bataticola: 473; 649
--- rolfsii: 164; 191; 314; 441; 941
Scolytus multistriatus: attempted to form brood galleries in recently trimmed, apparently healthy elm trees, first report, 476
Scotland: 1028
Seed: growth pouches used for root-knot nematode resistance tests, 482
--- borne: infection of Piricularia setariae as seedborne pathogen in Setaria italica in India, 138; two downy mildews of corn internally seedborne, 1010; fungi carried by corn seed and their importance in causing corn diseases in the U.A.R., 53
---: transmission: of arabis mosaic virus in lettuce, 883; of bacterial leaf blotch disease of sorghum in south India, 63
--- treatments: chloroneb used as overtreatment of mercury-treated cottonseed not so effective as captan-PCNB, 558; evaluation of seed treatment with chloroneb as cotton seedling disease control measure, 558; hot water seed treatment of rice, effective in eradicating inoculum of Xanthomonas oryzae, 882; streptomycin in addition to regular seed treatment showed no significant control of bean halo blight in field plots, 544
Serum: anti-, sheep-produced antibodies for use in detecting plant-pathogenic bacteria, 50
Silverleaf: Stereum purpureum on peach and apricot in Chile, first report of, 212
Smut: covered kernel, of sorghum, 21 varieties of sorghum screened for resistance in Iraq, 690
---: stripe, on 'Merion' Kentucky bluegrass, chemical control of stripe smut, 355
Snow mold: of winter wheat: three pathogens show no significant pathogenic specialization, 810; larger plants of all varieties tested survived infection better than smaller plants in controlled environmental trials, 815
Soil: soil-temperature control system employing air as the heat-transfer medium, 873; incidence of Pythium and Phytophthora species in forest soils in the southeastern United States, 929; moisture, and ozone related to occurrence of weather fleck on tobacco in Florida, 1966, 72
---: amendments, sawdust + balanced NPK fertilizers applied to soil suppressed root knot of okra and tomato and increased yield in India, 861; reduction of lesion nematodes in soybean meal and oil-amended soils, 1021
---: treatments: comparison of soil treatments for control of root rot of peas, 125; comparison of some methyl-containing amino acids and soil amendments for control of Aphanomyces root rot of peas, 125; effects of preplanting soil treatments on peanut losses and reduction of pod mycoflora in Georgia, 461; fumigation studies for control of nematodes of wheat in Nebraska resulted in good control, 637; fumigation with D-D gave good control of plant-parasitic nematodes in corn, however, did not result in significant increase in yields, 15; fumigants, for control of root rot of cotton, 331; soil fumigation by injection and irrigation for control of nematodes on hops, 794; nematocides in combination with fungicidal emergence stimulants may help reduce population of tobacco cyst nematodes in field, 609; in-furrow treatment with PCNB indicates some hope for control of head smut of corn, 626; liming of soil reduced incidence of Fusarium wilt (race 2) in tomato (variety Manapal), 645; plant-bed applications of 2,6-dichloro-4-nitroaniline effective in controlling sclerotial blight of sweetpotato, 191; control of southern blight on peanut, deep plowing and PCNB treatment of soil, and raised beds effective on irrigated fields in Texas, 314; sulfur, for sweetpotato soil rot control, 271
Solanum nigrum: nightshade, as suspect of bacterial spot, Xanthomonas vesicatoria, 659
Soot: damage to greenhouse plants, 712
Sorghum: mosaic disease, observed in four southern States, 200
---: bacterial leaf blotch, seed transmission, in India, 63
---: cotton root-knot nematode, associated with damage to grain sorghum in west Texas, 29
---: covered kernel smut, 21 varieties of sorghum screened for resistance in

Iraq, 690
---: Fusarium scab in Nigeria, 174
---: maize dwarf mosaic, threat to grain sorghum, 777
---: red stalk rot, relationship of root injury to cell death and spread of Colletotrichum graminicola in stalk tissue, 957
South Carolina: 557
South Dakota: 705; 981
Southern blight: of tomato, in Ghana, Molokai variety showed high resistance to infection in laboratory inoculations, 941
Soybean: brown stem rot incidence in Iowa, 438
---: cyst nematode: nature of resistance to, 771; first reported occurrence in Louisiana, 823
---: root and stem rot, caused by Rhizoctonia solani, 872
Spearmint: two Longidorus spp. found on, first report in Sudan, 30
Sphaceloma populi: poplar spotted anthracnose, first North American record of the pathogen in Oklahoma, 785
Sphacelotheca reiliana: 626
--- sorghi: 690
Sphaerotheca fuliginea: 1079
Spread: of rice blast in small fields, 243
Spruce (see Picea)
Stem-end rot: of mango, mode of infection and prevention, 744
Stemphylium botryosum: 741
Storage: cotton boll rot organisms stored on infected cotton bolls at least 8 years without loss in viability or virulence, 855; decay of crowns of crownvetch, captan promising as fungicidal dip for control, 820; effect of different drying temperatures and levels of relative humidity during storage on longevity of dried cultures of microorganisms, 1055; effect of dry storage versus subculturing on morphology and virulence of microorganisms, 1058; storage life influenced by dapple apple and stem pitting viruses, 293; wax emulsion additives for control of storage decay in lemons, 781
Strains: evidence that virus of corn and small grains in Ohio is strain of wheat streak mosaic virus, 207; predominating strains of barley yellow dwarf virus in New York showing changes during 10 year period, 195; 20 strains of Pellicularia sasakii detected and studied for first time in Taiwan, 682; of Pseudomonas phaseolicola, several bean varieties tested for reaction to, 20; transmission studies of the two strains of rice tungro virus, 877
Strawberry: angular leaf spot of, in Kentucky, 353; clone of Fragaria vesca (FV-72) from USSR more sensitive to strains of strawberry crinkle virus than indicator clones now generally in use, 288
Streptomyces ipomoea: 271
Stylosanthes gracilis: resistant or immune to undescribed species of Pratylenchus, 33
Subculture: effect of dry storage versus subculturing on morphology and virulence of microorganisms, 1058
Sudan: 30
Sugarbeet: plant pathogens cited as possible factor in unexpected preplant herbicide damage in sugarbeets, 86
---: viruses: curly top virus, possible incidence of in Iran, first record of, 976; in vitro increases in infectivity of cucumber mosaic virus in extracts from young sugarbeet leaves, 178
Sugarcane: leaf scald disease, isolates thought to be Xanthomonas albilineans from sugarcane infected with leaf scald disease proved to be pathogenic, 491; nematodes associated with plant disease, first report from Iraq, 809
---: maize dwarf mosaic virus, commercial varieties grown from cuttings can be symptomless hosts of MDMV, 761
Sulfur dioxide: causes plant damage in spring each year in New Jersey in form of foliar necrosis, 850
Sumac: leaf spot of skunkbush sumac seedlings controlled with captan or maneb, 700
Surgical treatments: for control of fusiform rust galls on loblolly pine, with greater care in trimming bark beyond upper and lower limits of gall surgery should have reasonable success, 505
Sweetgum: very susceptible to root rot caused by Corticium galactinum when pathogen was placed in contact with tap roots of tree seedlings, 500
Sweetpotato: sclerotial blight of, controlled by 2,6-dichloro-4-nitroaniline, 191; soil rot, development in Olivier slit loam as affected by lime or sulfur applications, 271
Sycamore: seedlings susceptible to root rot caused by Corticium galactinum, 500

Taiwan: 682
Tanzania: 276
Temperature: bacterial soft rot, bacterial necrosis and Alternaria rot of tomato

increased by commercial practice of field washing and postharvest chilling of mature-green fruit, 151; chilling-injury of potato tubers stored at 32°F in air, 1014; comparison of temperature effects on growth of Phymatotrichum omnivorum on solid media in laboratory, 926; constant temperature bath for pot-grown plants, 171; effect of different drying temperatures on longevity of dried cultures of microorganisms, 1055; hot water seed treatment for eradication of inoculum of Xanthomonas oryzae, 882; hot water treatments for control of endo- and ecto-parasitic nematodes in turf; 905; strong interaction between damping-off of small grain seedlings and temperature in Florida, 919; evidence that "needle curl" of loblolly pine seedlings may be caused by excessively high temperatures, 805; postharvest chilling of mature-green tomatoes increased incidence of bacterial soft rot, 151; influence on reproduction, male-female-juvenile ratio, and root to soil migration of Pratylenchus penetrans, 78; a soil-temperature control system employing air as the heat-transfer medium, 873; effects of soil temperature and plant age on severity of Cylindrocladium rot of peanut, 986; effect on sporulation of Phytophthora phaseoli on 'Fordhook' lima bean stems, 253; effect on symptom development of Lambert mottle of sweet cherries, 269; effect of temperature and hydrogen peroxide on extraction of burrowing nematodes from citrus roots, 1024; temperature appears to be one limiting factor in infection and pathogenesis of Phymatotrichum omnivorum on cotton, 926

Tennessee: 4; 455; 731

Texas: 133; 536; 777

Thailand: 960

Thielaviopsis basicola: 864

Tilletia caries: 287

Tip burn: of carnation, 793

Tobacco: burley, survey in North Carolina to determine viruses present, distribution, and relation of previous cropping history to occurrence, 409; occurrence of diseases in Kentucky in 1966, 866; ozone-sensitive plants used to detect presence of oxidants on the Delmarva Peninsula, 493; weather fleck, ozone and soil moisture related to occurrence in Florida, 1966, 72

---: black shank disease: effect of calcium nutrition on susceptibility of tobacco seedlings, 641; race 1 identified in five counties in Kentucky and three in Tennessee, 731

---: brown root rot, study to provide evidence for relationship between causes of brown root rot of tobacco and preceding crop, particularly soybean, 225

Tolyposporium penicillariae: 468

Tomato: amendments of sawdust + NPK fertilizer to soil suppressed root-knot infestation and increased yield of tomatoes in India, 861

---: bacterial canker, found for first time in Chile, 396

---: bacterial necrosis, Alternaria rot and bacterial soft rot increased by commercial practice of field washing and postharvest chilling of mature-green tomatoes, 151

---: bacterial spot, several species, varieties and plant introductions tested for reaction of foliage of Xanthomonas vesicatoria isolates, 25

---: Fusarium wilt, inhibition of Fusarium wilt (race 2) in tomato (variety Manapal) when field soil was amended with hydrated lime, 645

---: root knot of, nematocides superior to crop residues for control of, 219

---: southern blight: controlled by crop rotation or soil treatment with fumigant fungicides, 314; Molokai variety showed high resistance to infection by Sclerotium rolfsii, in laboratory inoculations in Ghana, 941

---: viruses: tomato aspermy virus: completely suppresses viable seed in tomato fruits, three chrysanthemum viruses similar but not identical to, 723; curly top virus in Iowa in 1967, 1069

Transmission: of Pseudomonas syringae in peach trees by bud propagation, 666

---: viruses: transmission of exocortis virus of citron plants due to contaminated tools, 410; of exocortis virus of citrus by dodder, grafting, and mechanical methods, 868; interaction of trichomes, abrasives, and mechanics of applying inoculum on transmission of viruses, 945; of two strains of rice tungro virus, 877; seed transmission of arabis mosaic virus in lettuce, 883; soilborne wheat mosaic virus best maintained for transmission by root washings, 1005; of stem pitting virus of apple, pattern spread in orchard suggests root grafting could account for most of natural transmission, 291

Trichoderma viride: 601
Trichodorus sp.: 905
Trichomes: virus transmission, interaction of trichomes, abrasives, and mechanics of applying inoculum on transmission, 945
Trimming: of healthy elm trees attracted colonizing smaller European elm bark beetles in Detroit, Michigan, 476
Tsuga: Phacidiopycnis pseudotsugae associated with bleeding cankers of, first report of fungus affecting naturally, 807
Tuberculina maxima: 89
Turkey: 893
Tylenchorhynchus sp.: 809
Tylenchulus semipenetrans: 317; 995
Typhula idahoensis: 810; 815
--- incarnata: 810

United Arab Republic: 53
United States: 182; Index to genera of pathogens listed in the "Index of Plant Diseases in the United States"--4 parts, 235, 323, 419, 515
Uromyces viciae-fabae: 595
Ustilago avenae: 844; 846
--- striiformis: 355

Vascular mycosis: on oak in Russia, possible relation to oak wilt in United States, 739
Vectors associated with: barley yellow dwarf virus in Iowa, greenbug Schizaphis graminum seemed principal vector, 527; barley yellow dwarf virus in South Dakota, Macrosiphum avenae, Rhopalosiphum spp. and Schizaphis graminum, 981; leaf-mottle-yellows disease in Philippines, psyllid Diaphorina citri, 692; virus-like disorder of rice in India transmitted by leafhopper Nephotettix bipunctatus, 300; rice yellow dwarf virus, virus experimentally transmitted by Nephotettix bipunctatus, 1040
Verticillium albo-atrum: 212; 449; 589; 608; 864
--- dahliae: 98; 608
--- nigrescens: 760
Vetch: (Vicia spp.), additional Vicia species found resistant to root-knot nematodes, 614
---: crown: storage decay of crowns controlled by captan fungicide dip treatment, 820
Vinca rosea: graft-transmissible disease causes flower phyllody, first report on transmission of yellows-type virus disease in North Africa, 416; twig blight, new disease in India caused by Colletotrichum sp., 480
Viola adunca: non-rosaceous host of Phragmidium (Uredinales), 766
Virginia: 169; 856
Virus: cross-transmission studies with Citrus and Prunus viruses, 64; detection in vegetatively propagated plant introductions under quarantine in the United States, 1957-1967, 715; nature and distribution of pome fruit viruses in Poland, 117; nature and distribution of stone fruit viruses in Poland, 115; of stone fruits, mechanically transmitted stone fruit viruses and their selective host range, 786
---: abaca mosaic, propagation of virus-free suckers from mosaic-infected abaca corms, 405
---: of apple: influence of dapple apple and stem pitting on tree growth and fruit yields and storage life, 293; stem pitting, pattern of spread in orchard suggests root grafting would account for most of natural transmission, 291; the name "flat apple virus" is proposed as the name for flat apple disease, 509; possibly a new virus disease of apple, graft-transmissible "red-ring" disease, 281
---: arabis mosaic virus, of lettuce in England, seed transmitted, 883
---: aspermy related viruses, from chrysanthemum pathogenic to tomato but not identical to tomato aspermy virus, 723
---: barley yellow dwarf: aluminum foil fails to protect winter oats from aphid vectors, 354; field infectivities of some aphid vectors of BYDV in South Dakota, 981; of oats, tolerance to, in several winter oats, 908; relationship of low-level jet winds with early cereal aphid and BYDV detection in Iowa, 527; report of predominating strains of BYDV in New York showing changes during 10-year period, 195; wild oat species show resistance in California test, 391
---: bean yellow mosaic, on broadbean crop in Iran, identified as one of three diseases causing reduced yields of over 50% in 1966, 595
---: western celery mosaic virus identified in celery from commercial plantings in Florida, 502
---: of cherry: apple chlorotic leaf spot virus (CLSV) and tobacco mosaic virus (TMV) isolated from cherry trees in New York indexing survey, 823; Lambert mottle, symptom development

affected by temperature, 269
---: of citrus: cachexia virus of citrus, Citrus macrophylla 'Alemow' used as rootstock for lemons in California susceptible to virus, 525; citrange stunt virus of citrus successfully transmitted by dodder, 629; exocortis virus of citron plants, disease transmitted mechanically by contaminated budding tools, 410; exocortis virus of citrus, successfully transmitted from citrus to herbaceous plants and back to citrus, 868; impietratura-diseased grapefruit fruits in Cyprus, effect of rootstock on incidence, 684; leaf-mottle-yellows disease of citrus in Philippines, may be virus responsible for severe tree losses in last 10 years, 692
---: corn: of corn and small grains in Ohio, thought to be strain of wheat streak mosaic virus, 207; corn stunt, reaction of dent corn hybrid to Mississippi isolate of CSV and to a mechanically transmissible mosaic virus isolate found on Johnson grass, 1044; nonpersistent virus found infecting corn in South Dakota, 705; positive relationship between virus ratings and survival of plants with ears, regardless of specific virus or viruses involved, 377
---: of cucumber, varietal performance of trellised cucumbers during serious outbreak of mosaic virus in western North Carolina, 431
---: cucumber mosaic: on Phlox, 768; from sugarbeet, in vitro increase in infectivity of virus, 178
---: curly top, on tomatoes and pumpkins in Iowa in 1967, 1069
---: of elm, Morsus ulmi (elm phloem necrosis), 511
---: fig mosaic: in Turkey, present in all nursery stock, 893
---: of iris: severe iris mosaic, symptomless occurrence in bulbous iris, 952; yellow mosaic in Iris, spread reduced by oil sprays in Israel, 318
---: on Johnson grass, plants with mosaic symptoms found in Louisiana, 108
---: on lettuce, tomato spotted wilt virus occurring naturally in Argentina, first report, 225
---: of oats, soilborne mosaic, resistance observed in 43 selections of Avena sterilis tested at Quincy, Florida, 120
---: maize dwarf mosaic: distribution in Iowa in 1966, 4 new locations, 388; additional data on distribution and properties on corn and Johnson grass, in Alabama, 385; in Texas, identification, transmission, and distribution of, on corn and grain sorghum, 777; recovered from commercial varieties of sugarcane, 761
---: of peach: maintaining virus-free cultivars of peach in Michigan, 261; peach wart, 'Shiro' plum a rapid greenhouse indicator of virus, 831; of Prunus spp., symptoms of peach wart virus in several Prunus spp. and peach and nectarine varieties, 42; peach yellow bud mosaic virus, transmission to peach, plum, and apricot seedlings by dagger nematode, 841; first report in Israel of bud-transmissible disease which resembles peach yellows, 733
---: Prunus ringspot virus and cherry yellows virus, incidence in Washington peach orchards possibly caused by infected rootstocks and budwood, 1063
---: stunt: of peanut: drastically reduced yield and quality of commercially grown peanuts, 856; stunt disease survey in Virginia, 1966, 169
---: of pear, mechanical transmission of pear viruses from pear to Chenopodium amaranticolor, 283
---: of pepper, identification of viruses affecting commercial varieties of pepper in New Jersey, 709
---: of periwinkle, first report of transmission of yellows-type virus disease in North Africa, 416
---: of potatoes, purple-top (aster yellows) in Guatemala, 265
---: of rice: inoculation with maize dwarf mosaic caused only localized infection, rice exposed to MDMV in field was not infected, 381; transmission of virus-like disease of rice by leafhopper, in India, 300; two strains of rice tungro virus identified and designated "S" and "M", 877; rice yellow dwarf, occurrence and transmission of in India, 1040
---: of sorghum, occurrence of a potentially destructive mosaic disease in four southern States, 200
---: strawberry crinkle virus, a more sensitive strawberry virus indicator clone of Fragaria vesca from the USSR, 288
---: of sugarbeet, possible incidence of curly top virus in Iran, first record of, 976
---: of tobacco, survey of virus diseases in North Carolina, 409
---: tobacco mosaic, natural infection of wisteria and cultivated elder in Rhode Island, apparently first report of, 1053

☆ U. S. GOVERNMENT PRINTING OFFICE : 1968 343-150/6

---: transmission, interaction of trichomes, abrasives and mechanics of applying inoculum on transmission of viruses, 945
---: of wheat: soilborne mosaic, maintenance of soilborne wheat mosaic virus cultures by transfer through root washings, 1005; wheat streak mosaic, first report from Iowa identifying causal virus and reporting recovery from both wheat and corn, 389

Washington: 94; 911; 1063
Watermelon (see Cucurbits)
Weather: relation of rainfall and temperature to late blight of potato in Greece, 75
Weather fleck: of cigar-wrapper tobacco, occurrence related to ozone and soil moisture in Florida, 1966, 72
West Virginia: 430; 1038
Wheat: bunt, wheat introductions PI 192383, PI 116301, and M63-1223, previously resistant to all known races of common bunt found susceptible to new race of Tilletia caries (designated X-1), 287
---: leaf blight, effectively controlled in India by fungicidal sprays, 950
---: nematodes: associated with roots of wheat, their population fluctuations and fumigation studies surveyed in Nebraska, 637; effect of variety and fumigation on nematode populations in Florida, 915
---: powdery mildew, 1966 epiphytotic in eastern Washington, 94
---: rust: leaf and stem: in Rumania, 14 races of leaf rust and four races of stem rust identified in 1964 and 1965, 358; control of by different formulations of 1,4-oxathiin derivatives, 336
---: ---: stem: races of Puccinia graminia tritici identified from Texas collections, 1926-1965, 536; chemically induced changes in urediospores of Puccinia graminis tritici, 904; minimum data apparently necessary for adequate prediction of field infection in experimental initiation of rust epidemics, 248
---: ---: stripe: induction of infection during summer in southeastern Washington, 911; comparison of root systems produced by healthy and stripe rust-inoculated wheat in mist-, water-, and sand-culture, 1074; single gene control of stripe rust resistance, 1041; effect on yield and its components of six winter wheat varieties, 702; stripe, leaf and stem rust, six winter wheats showed resistance to all three rusts in Rumanian tests in 1965, then tested against stem rust cultures in United States in 1966, 362
---: snow mold: of winter wheat, no evidence of pathogenic specialization among three pathogens, 810; of winter wheat, effect of plant size on resistance, 815
---: stand loss in Florida affected by temperature, seed treatment and soil fumigation, 919
---: virus: soilborne mosaic, maintenance of soilborne wheat mosaic virus cultures by transfer through root washings, 1005; streak mosaic virus, incidence in Iowa in 1966, 389
Wilt: branch, apparently a new disease of grape, widespread throughout grape-producing regions in Iraq, caused by Hendersonula toruloidea, 371
---: Fusarium: of celery, relationships of celery-wilt Fusarium to two races of pea-wilt, 888; of chrysanthemum, effect of Meloidogyne hapla and Fusarium oxysporum on severity of, 736; of coffee, new record in India, 758; of cumin, soil treatment with SMDC reduced incidence of fungus, 629; oxysporum lycopersici (race 2), liming of soil decreased incidence, 645
---: oak: antibiotic treatments for possible control in northern pin oaks, 160; mats and pressure cushions found on white oak for first time, West Virginia, 430; evaluation of small oak bark beetle as a vector, 755
---: Verticillium: first report of isolation of Verticillium nigrescens from cotton, in Missouri, 760; of peppermint, wilt effectively reduced in muck soils after 5 years' continuous cropping with corn or after reed canary grass sod, 449; resistance of 21 collections of noncultivated safflower species evaluated, 589
Wisconsin: 427
Wisteria: tobacco mosaic virus, apparently first report of natural infection of, 1053

Xanthomonas albilineans: 491; preservation of, 739
--- fragariae: 353
--- malvacearum: 1066
--- oryzae: 882
--- oryzicola: a proposed name change, 928
--- phaseoli: 622
--- pruni: 345; 351
--- rubrisorghi: 63
--- vesicatoria: 25; 57; 320; 497; 659; 841

UNITED STATES DEPARTMENT OF AGRICULTURE
AGRICULTURAL RESEARCH SERVICE
Crops Research Division
Beltsville, Maryland 20705

Postage and Fees Paid
U. S. DEPARTMENT OF AGRICULTURE

Official Business